

The Project Gutenberg EBook of Groups of Order p^m Which Contain Cyclic Subgroups of Order $p^{(m-3)}$, by Lewis Irving Neikirk

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you'll have to check the laws of the country where you are located before using this ebook.

Title: Groups of Order p^m Which Contain Cyclic Subgroups of Order $p^{(m-3)}$

Author: Lewis Irving Neikirk

Release Date: April 24, 2015 [EBook #9930]

Language: English

Character set encoding: ASCII

*** START OF THIS PROJECT GUTENBERG EBOOK GROUPS OF ORDER P^M WHICH ***

Produced by Cornell University, Joshua Hutchinson, Lee
Chew-Hung, John Hagerson, and the Online Distributed
Proofreading Team

TRANSCRIBER'S NOTE

Minor typographical corrections and presentational changes have been
made without comment. The L^AT_EX source file may be downloaded from

www.gutenberg.org/ebooks/9930.

GROUPS OF ORDER p^m WHICH CONTAIN
CYCLIC SUBGROUPS OF ORDER p^{m-3}

BY

LEWIS IRVING NEIKIRK

SOMETIME HARRISON RESEARCH FELLOW IN MATHEMATICS

1905

INTRODUCTORY NOTE.

This monograph was begun in 1902-3. Class I, Class II, Part I, and the self-conjugate groups of Class III, which contain all the groups with independent generators, formed the thesis which I presented to the Faculty of Philosophy of the University of Pennsylvania in June, 1903, in partial fulfillment of the requirements for the degree of Doctor of Philosophy.

The entire paper was rewritten and the other groups added while the author was Research Fellow in Mathematics at the University.

I wish to express here my appreciation of the opportunity for scientific research afforded by the Fellowships on the George Leib Harrison Foundation at the University of Pennsylvania.

I also wish to express my gratitude to Professor George H. Hallett for his kind assistance and advice in the preparation of this paper, and especially to express my indebtedness to Professor Edwin S. Crawley for his support and encouragement, without which this paper would have been impossible.

LEWIS I. NEIKIRK.

UNIVERSITY OF PENNSYLVANIA, *May, 1905.*

GROUPS OF ORDER p^m , WHICH CONTAIN CYCLIC SUBGROUPS OF ORDER $p^{(m-3)1}$

BY

LEWIS IRVING NEIKIRK

Introduction.

The groups of order p^m , which contain self-conjugate cyclic subgroups of orders p^{m-1} , and p^{m-2} respectively, have been determined by BURNSIDE,² and the number of groups of order p^m , which contain cyclic non-self-conjugate subgroups of order p^{m-2} has been given by MILLER.³

Although in the present state of the theory, the actual tabulation of all groups of order p^m is impracticable, it is of importance to carry the tabulation as far as may be possible. In this paper *all groups of order p^m* (p being an odd prime) *which contain cyclic subgroups of order p^{m-3} and none of higher order* are determined. The method of treatment used is entirely abstract in character and, in virtue of its nature, it is possible in each case to give explicitly the generational equations of these groups. They are divided into three classes, and it will be shown that these classes correspond to the three partitions: $(m-3, 3)$, $(m-3, 2, 1)$ and $(m-3, 1, 1, 1)$, of m .

We denote by G an abstract group G of order p^m containing operators of order p^{m-3} and no operator of order greater than p^{m-3} . Let P denote one of these operators of G of order p^{m-3} . The p^3 power of every operator in G is contained in the cyclic subgroup $\{P\}$, otherwise G would be of order greater than p^m . The complete division into classes is effected by the following assumptions:

- I. There is in G at least one operator Q_1 , such that $Q_1^{p^2}$ is not contained in $\{P\}$.
- II. The p^2 power of every operator in G is contained in $\{P\}$, and there is at least one operator Q_1 , such that Q_1^p is not contained in $\{P\}$.
- III. The p th power of every operator in G is contained in $\{P\}$.

¹Presented to the American Mathematical Society April 25, 1903.

²*Theory of Groups of a Finite Order*, pp. 75-81.

³Transactions, vol. 2 (1901), p. 259, and vol. 3 (1902), p. 383.

The number of groups for Class I, Class II, and Class III, together with the total number, are given in the table below:

	I	II ₁	II ₂	II ₃	II	III	Total
$p > 3$ $m > 8$	9	$20 + p$	$6 + 2p$	$6 + 2p$	$32 + 5p$	23	$64 + 5p$
$p > 3$ $m = 8$	8	$20 + p$	$6 + 2p$	$6 + 2p$	$32 + 5p$	23	$63 + 5p$
$p > 3$ $m = 7$	6	$20 + p$	$6 + 2p$	$6 + 2p$	$32 + 5p$	23	$61 + 5p$
$p = 3$ $m > 8$	9	23	12	12	47	16	72
$p = 3$ $m = 8$	8	23	12	12	47	16	71
$p = 3$ $m = 7$	6	23	12	12	47	16	69

Class I.

1. *General notations and relations.*—The group G is generated by the two operators P and Q_1 . For brevity we set⁴

$$Q_1^a P^b Q_1^c P^d \dots = [a, b, c, d, \dots].$$

Then the operators of G are given each uniquely in the form

$$[y, x] \left(\begin{array}{l} y = 0, 1, 2, \dots, p^3 - 1 \\ x = 0, 1, 2, \dots, p^{m-3} - 1 \end{array} \right).$$

We have the relation

$$Q_1^{p^3} = P^{hp^3}. \quad (1)$$

There is in G , a subgroup H_1 of order p^{m-2} , which contains $\{P\}$ self-conjugately.⁵ The subgroup H_1 is generated by P and some operator $Q_1^y P^x$ of G ; it then contains Q_1^y and is therefore generated by P and $Q_1^{p^2}$; it is also self-conjugate in $H_2 = \{Q_1^p, P\}$ of order p^{m-1} , and H_2 is self-conjugate in G .

From these considerations we have the equations⁶

$$Q_1^{-p^2} P Q_1^{p^2} = P^{1+kp^{m-4}}, \quad (2)$$

$$Q_1^{-p} P Q_1^p = Q_1^{\beta p^2} P^{\alpha_1}, \quad (3)$$

$$Q_1^{-1} P Q_1 = Q_1^{bp} P^{a_1}. \quad (4)$$

⁴With J. W. YOUNG, *On a certain group of isomorphisms*, American Journal of Mathematics, vol. 25 (1903), p. 206.

⁵BURNSIDE: *Theory of Groups*, Art. 54, p. 64.

⁶*Ibid.*, Art. 56, p. 66.

2. *Determination of H_1 . Derivation of a formula for $[yp^2, x]^s$.*—From (2), by repeated multiplication we obtain

$$[-p^2, x, p^2] = [0, x(1 + kp^{m-4})];$$

and by a continued use of this equation we have

$$[-yp^2, x, yp^2] = [0, x(1 + kp^{m-4})y] = [0, x(1 + kyp^{m-4})] \quad (m > 4)$$

and from this last equation,

$$[yp^2, x]^s = [syp^2, x\{s + k\binom{s}{2}yp^{m-4}\}]. \quad (5)$$

3. *Determination of H_2 . Derivation of a formula for $[yp, x]^s$.*—It follows from (3) and (5) that

$$[-p^2, 1, p^2] = \left[\beta \frac{\alpha_1^p - 1}{\alpha_1 - 1} p^2, \alpha_1^p \left\{ 1 + \frac{\beta k}{2} \frac{\alpha_1^p - 1}{\alpha_1 - 1} p^{m-4} \right\} \right] \quad (m > 4).$$

Hence, by (2),

$$\begin{aligned} \beta \frac{\alpha_1^p - 1}{\alpha_1 - 1} p^2 &\equiv 0 \pmod{p^3}, \\ \alpha_1^p \left\{ 1 + \frac{\beta k}{2} \frac{\alpha_1^p - 1}{\alpha_1 - 1} p^{m-4} \right\} + \beta \frac{\alpha_1^p - 1}{\alpha_1 - 1} hp^2 &\equiv 1 + kp^{m-4} \pmod{p^{m-3}}. \end{aligned}$$

From these congruences, we have for $m > 6$

$$\alpha_1^p \equiv 1 \pmod{p^3}, \quad \alpha_1 \equiv 1 \pmod{p^2},$$

and obtain, by setting

$$\alpha_1 = 1 + \alpha_2 p^2,$$

the congruence

$$\frac{(1 + \alpha_2 p^2)^p - 1}{\alpha_2 p^3} (\alpha_2 + h\beta) p^3 \equiv kp^{m-4} \pmod{p^{m-3}};$$

and so

$$(\alpha_2 + h\beta) p^3 \equiv 0 \pmod{p^{m-4}},$$

since

$$\frac{(1 + \alpha_2 p^2)^p - 1}{\alpha_2 p^3} \equiv 1 \pmod{p^2}.$$

From the last congruences

$$(\alpha_2 + h\beta)p^3 \equiv kp^{m-4} \pmod{p^{m-3}}. \quad (6)$$

Equation (3) is now replaced by

$$Q_1^{-p} P Q_1^{-p} = Q_1^{\beta p^2} P^{1+\alpha_2 p^2}. \quad (7)$$

From (7), (5), and (6)

$$[-yp, x, yp] = [\beta xyp^2, x\{1 + \alpha_2 yp^2\} + \beta k \binom{x}{2} yp^{m-4}].$$

A continued use of this equation gives

$$\begin{aligned} [yp, x]^s &= [syp + \beta \binom{s}{2} xyp^2, \\ &\quad xs + \binom{s}{2} \{\alpha_2 xyp^2 + \beta k \binom{x}{2} yp^{m-4}\} + \beta k \binom{s}{3} x^2 yp^{m-4}]. \end{aligned} \quad (8)$$

4. *Determination of G.*—From (4) and (8),

$$[-p, 1, p] = [Np, a_1^p + Mp^2].$$

From the above equation and (7),

$$a_1^p \equiv 1 \pmod{p^2}, \quad a_1 \equiv 1 \pmod{p}.$$

Set $a_1 = 1 + a_2p$ and equation (4) becomes

$$Q_1^{-1} P Q_1 = Q_1^{bp} P^{1+a_2p}. \quad (9)$$

From (9), (8) and (6)

$$[-p^2, 1, p^2] = \left[\frac{(1 + a_2p)^{p^2} - 1}{a_2p} bp, (1 + a_2p)^{p^2} \right],$$

and from (1) and (2)

$$\begin{aligned} \frac{(1 + a_2p)^{p^2} - 1}{a_2p} bp &\equiv 0 \pmod{p^3}, \\ (1 + a_2p)^{p^2} + bh \frac{(1 + a_2p)^{p^2} - 1}{a_2p} p &\equiv 1 + kp^{m-4} \pmod{p^{m-3}}. \end{aligned}$$

By a reduction similar to that used before,

$$(a_2 + bh)p^3 \equiv kp^{m-4} \pmod{p^{m-3}}. \quad (10)$$

The groups in this class are completely defined by (9), (1) and (10).

These defining relations may be presented in simpler form by a suitable choice of the second generator Q_1 . From (9), (6), (8) and (10)

$$[1, x]^{p^3} = [p^3, xp^3] = [0, (x+h)p^3] \quad (m > 6),$$

and, if x be so chosen that

$$x + h \equiv 0 \pmod{p^{m-6}},$$

$Q_1 P^x$ is an operator of order p^3 whose p^2 power is not contained in $\{P\}$. Let $Q_1 P^x = Q$. The group G is generated by Q and P , where

$$Q^{p^3} = 1, \quad P^{p^{m-3}} = 1.$$

Placing $h = 0$ in (6) and (10) we find

$$\alpha_2 p^3 \equiv a_2 p^3 \equiv kp^{m-4} \pmod{p^{m-3}}.$$

Let $\alpha_2 = \alpha p^{m-7}$, and $a_2 = ap^{m-7}$. Equations (7) and (9) are now replaced by

$$\begin{aligned} Q^{-p} P Q^p &= Q^{\beta p^2} P^{1+\alpha p^{m-5}}, \\ Q^{-1} P Q &= Q^{bp} P^{1+ap^{m-6}}. \end{aligned} \quad (11)$$

As a direct result of the foregoing relations, the groups in this class correspond to the partition $(m-3, 3)$. From (11) we find⁷

$$[-y, 1, y] = [byp, 1 + ayp^{m-6}] \quad (m > 8).$$

It is important to notice that by placing $y = p$ and p^2 in the preceding equation we find that⁸

$$b \equiv \beta \pmod{p}, \quad a \equiv \alpha \equiv k \pmod{p^3} \quad (m > 7).$$

A combination of the last equation with (8) yields⁹

$$\begin{aligned} [-y, x, y] &= [bxyp + b^2 \binom{x}{2} yp^2, \\ &x(1 + ayp^{m-6}) + ab \binom{x}{2} yp^{m-5} + ab^2 \binom{x}{3} yp^{m-4}] \quad (m > 8). \end{aligned} \quad (12)$$

⁷For $m = 8$ it is necessary to add $a^2 \binom{y}{2} p^4$ to the exponent of P and for $m = 7$ the terms $a(a + \frac{abp}{2}) \binom{y}{2} p^2 + a^3 \binom{y}{3} p^3$ to the exponent of P , and the term $ab \binom{y}{2} p^2$ to the exponent of Q . The extra term $27ab^2k \binom{y}{3}$ is to be added to the exponent of P for $m = 7$ and $p = 3$.

⁸For $m = 7$, $ap^2 - \frac{a^2 p^3}{2} \equiv ap^2 \pmod{p^4}$, $ap^3 \equiv kp^3 \pmod{p^4}$. For $m = 7$ and $p = 3$ the first of the above congruences has the extra terms $27(a^3 + ab\beta k)$ on the left side.

⁹For $m = 8$ it is necessary to add the term $a \binom{y}{2} xp^4$ to the exponent of P , and for $m = 7$ the terms $x\{a(a + \frac{abp}{2}) \binom{y}{2} p^2 + a^3 \binom{y}{3} p^3\}$ to the exponent of P , with the extra term $27ab^2k \binom{y}{3} x$ for $p = 3$, and the term $ab \binom{y}{2} xp^2$ to the exponent of Q .

From (12) we get¹⁰

$$[y, x]^s = [ys + by\{(x + b\binom{x}{2}p)\binom{s}{2} + x\binom{s}{3}p\}p, \\ xs + ay\{(x + b\binom{x}{2}p + b^2\binom{x}{3}p^2)\binom{s}{2} \\ + (bx^2p + 2b^2x\binom{x}{2}p^2)\binom{s}{3} + bx^2\binom{s}{4}p^2\}p^{m-6}] \quad (m > 8). \quad (13)$$

5. *Transformation of the Groups.*—The general group G of Class I is specified, in accordance with the relations (2) (11) by two integers a, b which (see (11)) are to be taken mod p^3 , mod p^2 , respectively. Accordingly setting

$$a = a_1p^\lambda, \quad b = b_1p^\mu,$$

where

$$dv[a_1, p] = 1, \quad dv[b_1, p] = 1 \quad (\lambda = 0, 1, 2, 3; \mu = 0, 1, 2),$$

we have for the group $G = G(a, b) = G(a, b)(P, Q)$ the generational determination:

$$G(a, b) : \begin{cases} Q^{-1} P Q = Q^{b_1p^{\mu+1}} P^{1+a_1p^{m+\lambda-6}} \\ Q^{p^3} = 1, \quad P^{p^{m-3}} = 1. \end{cases}$$

Not all of these groups however are distinct. Suppose that

$$G(a, b)(P, Q) \sim G(a', b')(P', Q'),$$

by the correspondence

$$C = \begin{bmatrix} Q, & P \\ Q'_1, & P'_1 \end{bmatrix},$$

where

$$Q'_1 = Q'^y P'^x p^{m-6}, \quad \text{and} \quad P'_1 = Q'^y P'^x,$$

¹⁰For $m = 8$ it is necessary to add the term $\frac{1}{2}axy\binom{s}{2}[\frac{1}{3}y(2s-1) - 1]p^4$ to the exponent of P , and for $m = 7$ the terms

$$x\left\{\frac{a}{2}\left(a + \frac{ab}{2}\right)\left(\frac{2s-1}{3}y - 1\right)\binom{s}{2}yp^2 + \frac{a^3}{3!}\left(\binom{s}{2}y^2 - (2s-1)y + 2\right)yp^3 \right. \\ \left. + \frac{a^2bxy^2}{2}\binom{s}{3}\frac{3s-1}{2}p^3 + \frac{a^2b}{2}\left(\frac{s(s-1)^2(s-4)}{4!}y - \binom{s}{3}\right)yp^3\right\}$$

with the extra terms

$$27abxy\left\{\frac{bk}{3!}\left[\binom{s}{2}y^2 - (2s-1)y + 2\right]\binom{s}{3} + x(b^2k + a^2)(2y^2 + 1)\binom{s}{3}\right\},$$

for $p = 3$, to the exponent of P , and the terms $\frac{ab}{2}\{2s - \frac{1}{3}y - 1\}\binom{s}{2}xyp^2$ to the exponent of Q .

with y' and x prime to p .

Since

$$Q^{-1} P Q = Q^{bp} P^{1+ap^{m-6}},$$

then

$$Q_1'^{-1} P_1' Q_1' = Q_1'^{bp} P_1'^{1+ap^{m-6}},$$

or in terms of Q' , and P'

$$\begin{aligned} [y + b'xy'p + b'^2 \binom{x}{2} y' p^2, x(1 + a'y'p^{m-6}) + a'b' \binom{x}{2} y' p^{m-5} \\ + a'b'^2 \binom{x}{3} y' p^{m-4}] = [y + by'p, x + (ax + bx'p)p^{m-6}] \quad (m > 8) \end{aligned}$$

and

$$by' \equiv b'xy' + b'^2 \binom{x}{2} y' p \pmod{p^2}, \quad (14)$$

$$ax + bx'p \equiv a'y'x + a'b' \binom{x}{2} y' p + a'b'^2 \binom{x}{3} y' p^2 \pmod{p^3}. \quad (15)$$

The necessary and sufficient condition for the simple isomorphism of these two groups $G(a, b)$ and $G(a', b')$ is, that the above congruences shall be consistent and admit of solution for x, y, x' and y' . The congruences may be written

$$\begin{aligned} b_1 p^\mu &\equiv b'_1 x p^{\mu'} + b_1'^2 \binom{x}{2} p^{2\mu'+1} \pmod{p^2}, \\ a_1 x p^\lambda + b_1 x' p^{\mu+1} &\equiv \\ y' \{ a'_1 x p^{\lambda'} + a'_1 b'_1 \binom{x}{2} p^{\lambda'+\mu'+1} + a'_1 b_1'^2 \binom{x}{3} p^{\lambda'+2\mu'+2} \} &\pmod{p^3}. \end{aligned}$$

Since $dv[x, p] = 1$ the first congruence gives $\mu = \mu'$ and x may always be so chosen that $b_1 = 1$.

We may choose y' in the second congruence so that $\lambda = \lambda'$ and $a_1 = 1$ except for the cases $\lambda' \geq \mu + 1 = \mu' + 1$ when we will so choose x' that $\lambda = 3$.

The type groups of Class I for $m > 8$ ¹¹ are then given by

$$\begin{aligned} G(p^\lambda, p^\mu) : Q^{-1} P Q = Q^{p^{1+\mu}} P^{1+p^{m-6+\lambda}}, Q^{p^3} = 1, P^{p^{m-3}} = 1 \\ \left(\begin{array}{l} \mu = 0, 1, 2; \lambda = 0, 1, 2; \lambda \geq \mu; \\ \mu = 0, 1, 2; \lambda = 3 \end{array} \right). \quad (I) \end{aligned}$$

Of the above groups $G(p^\lambda, p^\mu)$ the groups for $\mu = 2$ have the cyclic subgroup $\{P\}$ self-conjugate, while the group $G(p^3, p^2)$ is the abelian group of type $(m-3, 3)$.

¹¹For $m = 8$ the additional term ayp appears on the left side of the congruence (14) and $G(1, p^2)$ and $G(1, p)$ become simply isomorphic. The extra terms appearing in congruence (15) do not effect the result. For $m = 7$ the additional term ay appears on the left side of (14) and $G(1, 1)$, $G(1, p)$, and $G(l, p^2)$ become simply isomorphic, also $G(p, p)$ and $G(p, p^2)$.

Class II.

1. General relations.

There is in G an operator Q_1 such that $Q_1^{p^2}$ is contained in $\{P\}$ while Q_1^p is not.

$$Q_1^{p^2} = P^{hp^2}. \quad (1)$$

The operators Q_1 and P either generate a subgroup H_2 of order p^{m-1} , or the entire group G .

Section 1.

2. Groups with independent generators.

Consider the first possibility in the above paragraph. There is in H_2 , a subgroup H_1 of order p^{m-2} , which contains $\{P\}$ self-conjugately.¹² H_1 is generated by Q_1^p and P . H_2 contains H_1 self-conjugately and is itself self-conjugate in G . From these considerations¹³

$$Q_1^{-p} P Q_1^p = P^{1+kp^{m-4}}, \quad (2)$$

$$Q_1^{-1} P Q = Q_1^{\beta p} P^{\alpha_1}. \quad (3)$$

3. Determination of H_1 and H_2 .

From (2) we obtain

$$[yp, x]^s = [syp, x\{s + k\binom{s}{2}yp^{m-4}\}] \quad (m > 4), \quad (4)$$

and from (3) and (4)

$$[-p, 1, p] = \left[\frac{\alpha_1^p - 1}{\alpha_1 - 1} \beta p, \alpha_1^p \left\{ 1 + \frac{\beta k}{2} \frac{\alpha_1^p - 1}{\alpha_1 - 1} p^{m-4} \right\} \right].$$

A comparison of the above equation with (2) shows that

$$\begin{aligned} \frac{\alpha_1^p - 1}{\alpha_1 - 1} \beta p &\equiv 0 \pmod{p^2}, \\ \alpha_1^p \left\{ 1 + \frac{\beta k}{2} \frac{\alpha_1^p - 1}{\alpha_1 - 1} p^{m-4} \right\} + \frac{\alpha_1^p - 1}{\alpha_1 - 1} \beta h p &\equiv 1 + kp^{m-4} \pmod{p^{m-3}}, \end{aligned}$$

and in turn

$$\alpha_1^p \equiv 1 \pmod{p^2}, \quad \alpha_1 \equiv 1 \pmod{p} \quad (m > 5).$$

Placing $\alpha_1 = 1 + \alpha_2 p$ in the second congruence, we obtain as in Class I

$$(\alpha_2 + \beta h)p^2 \equiv kp^{m-4} \pmod{p^{m-3}} \quad (m > 5). \quad (5)$$

¹²BURNSIDE, *Theory of Groups*, Art. 54, p. 64.

¹³*Ibid.*, Art. 56, p. 66.

Equation (3) now becomes

$$Q_1^{-1} P Q_1 = Q^\beta P^{1+\alpha_2 p}. \quad (6)$$

The generational equations of H_2 will be simplified by using an operator of order p^2 in place of Q_1 .

From (5), (6) and (4)

$$[y, x]^s = [sy + U_s p, sx + W_s p]$$

in which

$$\begin{aligned} U_s &= \beta \binom{s}{2} xy, \\ W_s &= \alpha_2 \binom{s}{2} xy + \left\{ \beta k \left[\binom{s}{2} \binom{x}{2} + \binom{s}{3} x^2 y \right] \right. \\ &\quad \left. + \frac{1}{2} \alpha k \left[\frac{1}{3!} s(s-1)(2s-1)y^2 - \binom{s}{2} y \right] x \right\} p^{m-5}. \end{aligned}$$

Placing $s = p^2$ and $y = 1$ in the above

$$[Q_1 P^x]^{p^2} = Q_1^{p^2} P^{xp^2} = P^{(x+h)p^2}.$$

If x be so chosen that

$$(x+h) \equiv 0 \pmod{p^{m-5}} \quad (m > 5)$$

$Q_1 P^x$ will be the required Q of order p^2 .

Placing $h = 0$ in congruence (5) we find

$$\alpha_2 p^2 \equiv k p^{m-4} \pmod{p^{m-3}}.$$

Let $\alpha_2 = \alpha p^{m-6}$. H_2 is then generated by

$$Q^{p^2} = 1, \quad P^{p^{m-3}} = 1.$$

$$Q^{-1} P Q = Q^{\beta p} P^{1+\alpha p^{m-5}}. \quad (7)$$

Two of the preceding formulæ now become

$$[-y, x, y] = [\beta x y p, x(1 + \alpha y p^{m-5}) + \beta k \binom{x}{2} y p^{m-4}], \quad (8)$$

$$[y, x]^s = [sy + U_s p, xs + W_s p^{m-5}], \quad (9)$$

where

$$U_s = \beta \binom{s}{2} xy$$

and¹⁴

$$W_s = \alpha \binom{s}{2} xy + \beta k \left\{ \binom{s}{2} \binom{x}{2} + \binom{s}{3} x^2 \right\} y p \quad (m > 6).$$

¹⁴For $m = 6$ it is necessary to add the terms $\frac{\alpha k}{2} \left\{ \frac{s(s-1)(2s-1)}{3!} y^2 - \binom{s}{2} y \right\} p$ to W_s .

4. *Determination of G.*

Let R_1 be an operation of G not in H_2 . R_1^p is in H_2 . Let

$$R_1^p = Q^{\lambda p} P^{\mu p}. \quad (10)$$

Denoting $R_1^a Q^b P^c R_1^d Q^e P^f \dots$ by the symbol $[a, b, c, d, e, f, \dots]$, all the operations of G are contained in the set $[z, y, x]$; $z = 0, 1, 2, \dots, p-1$; $y = 0, 1, 2, \dots, p^2-1$; $x = 0, 1, 2, \dots, p^{m-3}-1$.

The subgroup H_2 is self-conjugate in G . From this¹⁵

$$R_1^{-1} P R_1 = Q^{b_1} P^{a_1}, \quad (11)$$

$$R_1^{-1} Q R_1 = Q^{d_1} P^{c_1 p^{m-5}}. \quad (12)$$

In order to ascertain the forms of the constants in (11) and (12) we obtain from (12), (11), and (9)

$$[-p, 1, 0, p] = [0, d_1^p + Mp, Np^{m-5}].$$

By (10) and (8)

$$R_1^p Q R_1^p = P^{-\mu p} Q P^{\mu p} = Q P^{-a\mu p^{m-4}}.$$

From these equations we obtain

$$d_1^p \equiv 1 \pmod{p} \quad \text{and} \quad d_1 \equiv 1 \pmod{p}.$$

Let $d_1 = 1 + dp$. Equation (12) is replaced by

$$R_1^{-1} Q R_1 = Q^{1+dp} P^{e_1 p^{m-5}}. \quad (13)$$

From (11), (13) and (9)

$$[-p, 0, 1, p] = \left[\frac{a_1^p - 1}{a_1 - 1} b_1 + Kp, a_1^p + b_1 L p^{m-5} \right]$$

in which

$$K = a_1 b_1 \beta \sum_1^{p-1} \binom{a_1^y}{2}.$$

By (10) and (8)

$$R_1^{-p} P R_1^p = Q^{-\lambda p} P Q^{\lambda p} = P^{1+a\lambda p^{m-4}},$$

¹⁵BURNSIDE, *Theory of Groups*, Art. 24, p. 27.

and from the last two equations

$$a_1^p \equiv 1 \pmod{p^{m-5}}$$

and

$$a_1 \equiv 1 \pmod{p^{m-6}} \quad (m > 6); \quad a_1 \equiv 1 \pmod{p} \quad (m = 6).$$

$$\text{Placing } a_1 = 1 + a_2 p^{m-6} \quad (m > 6); \quad a_1 = 1 + a_2 p \quad (m = 6).$$

$$K \equiv 0 \pmod{p},$$

and¹⁶

$$\frac{a_1^p - 1}{a_1 - 1} b_1 \equiv b_1 p \equiv 0 \pmod{p^2}, \quad b_1 \equiv 0 \pmod{p}.$$

Let $b_1 = bp$ and we find

$$a_1^p \equiv 1 \pmod{p^{m-4}}, \quad a_1 \equiv 1 \pmod{p^{m-5}}.$$

Let $a_1 = 1 + a_3 p^{m-5}$ and equation (11) is replaced by

$$R_1^{-1} P R_1 = Q^{bp} P^{1+a_3 p^{m-5}}. \quad (14)$$

The preceding relations will be simplified by taking for R_1 an operator of order p . This will be effected by two transformations.

From (14), (9) and (13)¹⁷

$$[1, y]^p = \left[p, yp, \frac{-c_1 y}{2} p^{m-4} \right] = \left[0, (\lambda + y)p, \mu p - \frac{c_1 y}{2} p^{m-4} \right],$$

and if y be so chosen that

$$\lambda + y \equiv 0 \pmod{p},$$

$R_2 = R_1 Q^y$ is an operator such that R_2^p is in $\{P\}$.

Let

$$R_2^p = P^{lp}.$$

Using R_2 in the place of R_1 , from (15), (9) and (14)

$$[1, 0, x]^p = \left[p, 0, xp + \frac{ax}{2} p^{m-4} \right] = \left[0, 0, (x + l)p + \frac{ax}{2} p^{m-4} \right],$$

¹⁶ K has an extra term for $m = 6$ and $p = 3$, which reduces to $3b_1 c_1$. This does not affect the reasoning except for $c_1 = 2$. In this case change P^2 to P and c_1 becomes 1.

¹⁷The extra terms appearing in the exponent of P for $m = 6$ do not alter the result.

and if x be so chosen that

$$x + l + \frac{ax}{2}p^{m-5} \equiv 0 \pmod{p^{m-4}},$$

then $R = R_2P^x$ is the required operator of order p .

$R^p = 1$ is permutable with both Q and P . Preceding equations now assume the final forms

$$Q^{-1} P Q = Q^{\beta p} P^{1+ap^{m-5}}, \quad (15)$$

$$R^{-1} P R = Q^{bp} P^{1+ap^{m-4}}, \quad (16)$$

$$R^{-1} Q R = Q^{1+dp} P^{cp^{m-4}}, \quad (17)$$

with $R^p = 1$, $Q^{p^2} = 1$, $P^{p^{m-3}} = 1$.

The following derived equations are necessary¹⁸

$$[0, -y, x, 0, y] = [0, \beta xyp, x(1 + \alpha yp^{m-5}) + \alpha\beta \binom{x}{2} yp^{m-4}], \quad (18)$$

$$[-y, 0, x, -y] = [0, b xyp, x(1 + ayp^{m-4}) + ab \binom{x}{2} yp^{m-4}], \quad (19)$$

$$[-y, x, 0, y] = [0, x(1 + dyp), c xyp^{m-4}]. \quad (20)$$

From a consideration of (18), (19) and (20) we arrive at the expression for a power of a general operator of G .

$$[z, y, x]^s = [sz, sy + U_s p, sx + V_s p^{m-5}], \quad (21)$$

where¹⁹

$$\begin{aligned} U_s &= \binom{s}{2} \{bxz + \beta xy + dyz\}, \\ V_s &= \binom{s}{2} \left\{ \alpha xy + [axz + \alpha\beta \binom{x}{2} y + cyz + ab \binom{x}{2} z] p \right\} \\ &\quad + \alpha \binom{s}{3} \{bxz + \beta xy + dyz\} xp. \end{aligned}$$

5. *Transformation of the groups.* All groups of this section are given by equations (15), (16), and (17) with $a, b, \beta, c, d = 0, 1, 2, \dots, p-1$, and $\alpha = 0, 1, 2, \dots, p^2-1$, independently. Not all these groups, however, are distinct. Suppose that G and G' of the above set are simply isomorphic and that the correspondence is given by

$$C = \begin{bmatrix} R, & Q, & P \\ R'_1, & Q'_1, & P'_1 \end{bmatrix},$$

in which

$$\begin{aligned} R'_1 &= R'^{z''} Q'^{y''} P'^{x''} p^{m-4}, \\ Q'_1 &= R'^{z'} Q'^{y'} P'^{x'} p^{m-5}, \\ P'_1 &= R'^z Q'^y P'^x, \end{aligned}$$

¹⁸For $m = 6$ the term $a^2 \binom{x}{2} xp^2$ must be added to the exponent of P in (18).

¹⁹When $m = 6$ the following terms are to be added to V_s : $\frac{a^2 x}{2} \left\{ \frac{s(s-1)(2s-1)}{3!} y^2 - \binom{s}{2} y \right\} p$.

where x , y' and z'' are prime to p .

The operators R'_1 , Q'_1 , and P'_1 must be independent since R , Q , and P are, and that this is true is easily verified. The lowest power of Q'_1 in $\{P'_1\}$ is $Q'^{p^2}_1 = 1$ and the lowest power of R'_1 in $\{Q'_1, P'_1\}$ is $R'^p_1 = 1$. Let $Q'^{s'}_1 = P'^{sp^{m-5}}_1$.

This in terms of R' , Q' , and P' is

$$\left[s'z', y' \{ s' + d' \binom{s'}{2} z'p \}, s'x'p^{m-5} + c' \binom{s'}{2} y'z'p^{m-4} \right] = [0, 0, sxp^{m-5}].$$

From this equation s' is determined by

$$\begin{aligned} s'z' &\equiv 0 \pmod{p} \\ y' \{ s' + d' \binom{s'}{2} z'p \} &\equiv 0 \pmod{p^2}, \end{aligned}$$

which give

$$s'y' \equiv 0 \pmod{p^2}.$$

Since y' is prime to p

$$s' \equiv 0 \pmod{p^2}$$

and the lowest power of Q'_1 contained in $\{P'_1\}$ is $Q'^{p^2}_1 = 1$.

Denoting by $R'^{s''}_1$ the lowest power of R'_1 contained in $\{Q'_1, P'_1\}$.

$$R'^{s''}_1 = Q'^{s'p}_1 P'^{sp^{m-4}}_1.$$

This becomes in terms of R' , Q' , and P'

$$[s''z'', s''y''p, s''x''p^{m-4}] = [0, s'y'p, \{s'x' + sx\}p^{m-4}].$$

s'' is now determined by

$$s''z'' \equiv 0 \pmod{p}$$

and since z'' is prime to p

$$s'' \equiv 0 \pmod{p}.$$

The lowest power of R'_1 contained in $\{Q'_1, P'_1\}$ is therefore $R'^p_1 = 1$.

Since R , Q , and P satisfy equations (15), (16), and (17) R'_1 , Q'_1 , and P'_1 also satisfy them. Substituting in these equations the values of R'_1 , Q'_1 , and P'_1 and reducing we have in terms of R' , Q' , and P'

$$[z, y + \theta_1p, x + \phi_1p^{m-5}] = [z, y + \beta y'p, x(1 + \alpha p^{m-5}) + \beta xp^{m-4}], \quad (22)$$

$$[z, y + \theta_2p, x + \phi_2p^{m-4}] = [z, y + \beta y'p, x(1 + \alpha p^{m-4}) + \beta x'p^{m-4}], \quad (23)$$

$$[z', y' + \theta_3p, (x' + \phi_3p)p^{m-5}] = [z', y'(1 + dp), x(1 + dp)p^{m-5} + c xp^{m-4}], \quad (24)$$

in which

$$\begin{aligned}
\theta_1 &= d'(yz' - y'z) + x(b'z' + \beta'y'), \\
\theta_2 &= d'yz'' + b'xz'', \\
\theta_3 &= d'y'z'', \\
\phi_1 &= \alpha'xy' + \{\alpha'(\beta'y' + b'z')\binom{x}{2} + a'xz + c'(yz' - y'z)\}p, \\
\phi_2 &= \alpha'xy'' + a'xz'' + \alpha'b'\binom{x}{2}z'' + c'yz'', \\
\phi_3 &= c'yz''.
\end{aligned}$$

A comparison of the members of the above equations give six congruences between the primed and unprimed constants and the nine indeterminates.

$$\theta_1 \equiv \beta y' \pmod{p}, \quad (\text{I})$$

$$\phi_1 \equiv \alpha x + \beta x' p \pmod{p^2}, \quad (\text{II})$$

$$\theta_2 \equiv b y' \pmod{p}, \quad (\text{III})$$

$$\phi_2 \equiv a x + b x' \pmod{p}, \quad (\text{IV})$$

$$\theta_3 \equiv d y' \pmod{p}, \quad (\text{V})$$

$$\phi_3 \equiv c x + d x' \pmod{p}. \quad (\text{VI})$$

The necessary and sufficient condition for the simple isomorphism of the two groups G and G' is, *that the above congruences shall be consistent and admit of solution for the nine indeterminates, with the condition that x , y' and z'' be prime to p .*

For convenience in the discussion of these congruences, the groups are divided into six sets, and each set is subdivided into 16 cases.

The group G' is taken from the simplest case, and we associate with this case all cases, which contain a group G , simply isomorphic with G' . Then a single group G , in the selected case, simply isomorphic with G' , is chosen as a type.

G' is then taken from the simplest of the remaining cases and we proceed as above until all the cases are exhausted.

Let $\kappa = \kappa_1 p^{\kappa_2}$, and $dv_1[\kappa_1, p] = 1$ ($\kappa = a, b, \alpha, \beta, c$, and d).

The six sets are given in the table below.

I.

	α_2	d_2		α_2	d_2
A	0	0	D	2	0
B	0	1	E	1	1
C	1	0	F	2	1

The subdivision into cases and the results are given in Table II.

II.

	a_2	b_2	β_2	c_2	A	B	C	D	E	F
1	1	1	1	1						
2	0	1	1	1	A_1	B_1		C_2		E_2
3	1	0	1	1	A_1		C_1	D_1		
4	1	1	0	1	A_1		C_1	D_1		E_4
5	1	1	1	0	A_1		C_1	D_1		E_5
6	0	0	1	1	A_1	B_3	C_2	C_2	E_3	F_3
7	0	1	0	1	A_1	B_4	C_2	C_2		E_7
8	0	1	1	0	A_1	B_5	C_2	C_2	E_5	E_5
9	1	0	0	1	A_1	B_3	C_1	D_1	E_3	F_3
10	1	0	1	0	A_1		C_2	C_2		E_{10}
11	1	1	0	0	A_1		*	C_1		E_{11}
12	0	0	0	1	A_1	B_3	C_2	C_2	*	E_3
13	0	0	1	0	A_1	B_{10}	*	*	E_{10}	E_{10}
14	0	1	0	0	A_1	B_{11}	C_2	C_2	E_{11}	E_{11}
15	1	0	0	0	A_1	B_{10}	C_2	C_2	E_{10}	E_{10}
16	0	0	0	0	A_1	B_{10}	*	*	E_{10}	E_{10}

The groups marked (*) divide into two or three parts.

Let $ad - bc = \theta_1 p^{\theta_2}$, $\alpha_1 d - \beta c = \phi_1 p^{\phi_2}$ and $\alpha_1 b - a\beta = \chi_1 p^{\chi_2}$ with θ_1, ϕ_1 , and χ_1 prime to p .

III.

*	θ_2	ϕ_2	χ_2		*	θ_2	ϕ_2	χ_2	
C_{11}		1		D_1	D_{13}	1			D_1
C_{11}		0		C_1	D_{13}	0			C_2
C_{13}	1			C_1	D_{16}	1			C_1
C_{13}	0			C_2	D_{16}	0			C_2
C_{16}	1	1		D_1	E_{12}			1	F_3
C_{16}	1	0		C_1	E_{12}			0	E_3
C_{16}	0			C_2					

6. Types.

The type groups are given by equations (15), (16) and (17) with the values of the constants given in Table IV.

IV.

	a	b	α	β	c	d		a	b	α	β	c	d
A_1	0	0	1	0	0	1	E_1	0	0	p	0	0	0
B_1	0	0	1	0	0	0	E_2	1	0	p	0	0	0
B_3	0	1	1	0	0	0	E_3	0	1	p	0	0	0
B_4	0	0	1	1	0	0	E_4	0	0	p	1	0	0
B_5	0	0	1	0	1	0	E_5	0	0	p	0	1	0
B_{10}	0	1	1	0	κ	0	E_7	1	0	p	1	0	0
B_{11}	0	0	1	1	1	0	E_{10}	0	1	p	0	κ	0
C_1	0	0	p	0	0	1	E_{11}	0	0	p	1	1	0
C_2	ω	0	p	0	0	1	F_1	0	0	0	0	0	0
D_1	0	0	0	0	0	1	F_3	0	1	0	0	0	0

$\kappa = 1$, and a non-residue $(\text{mod } p)$,

$\omega = 1, 2, \dots, p-1$.

The congruences for three of these cases are completely analyzed as illustrations of the methods used.

B_{10} .

The congruences for this case have the special forms.

$$b'xz' \equiv \beta y' \pmod{p}, \quad (\text{I})$$

$$\alpha' y' \equiv \alpha \pmod{p}, \quad (\text{II})$$

$$b'xz'' \equiv by' \pmod{p}, \quad (\text{III})$$

$$\alpha'xy'' + \alpha'b'\binom{x}{2}z'' + c'yz'' \equiv ax + bx' \pmod{p}, \quad (\text{IV})$$

$$d \equiv 0 \pmod{p}, \quad (\text{V})$$

$$c'y'z'' \equiv cx \pmod{p}. \quad (\text{VI})$$

Since z' is unrestricted (I) gives $\beta \equiv 0$ or $\not\equiv 0 \pmod{p}$.

From (II) since $y' \not\equiv 0, \alpha \not\equiv 0 \pmod{p}$.

From (III) since $x, y', z'' \not\equiv 0, b \not\equiv 0 \pmod{p}$.

In (IV) $b \not\equiv 0$ and x' is contained in this congruence alone, and, therefore, a may be taken $\equiv 0$ or $\not\equiv 0 \pmod{p}$.

(V) gives $d \equiv 0 \pmod{p}$ and (VI), $c \not\equiv 0 \pmod{p}$.

Elimination of y' between (III) and (VI) gives

$$b'c'z''^2 \equiv bc \pmod{p}$$

so that bc is a quadratic residue or non-residue (mod p) according as $b'c'$ is a residue or non-residue.

The types are given by placing $a = 0$, $b = 1$, $\alpha = 1$, $\beta = 0$, $c = \kappa$, and $d = 0$ where κ has the two values, 1 and a representative non-residue of p .

C_2 .

The congruences for this case are

$$d'(yz' - y'z) \equiv \beta y' \pmod{p}, \quad (\text{I})$$

$$\alpha_1' xy' + a'xz' \equiv \alpha_1 x + \beta x' \pmod{p}, \quad (\text{II})$$

$$d'yz'' \equiv by' \pmod{p}, \quad (\text{III})$$

$$a'xz'' \equiv ax + bx' \pmod{p}, \quad (\text{IV})$$

$$d'z'' \equiv d \pmod{p}, \quad (\text{V})$$

$$cx + dx' \equiv 0 \pmod{p}. \quad (\text{VI})$$

Since z appears in (I) alone, β can be either $\equiv 0$ or $\not\equiv 0 \pmod{p}$. (II) is linear in z' and, therefore, $\alpha \equiv 0$ or $\not\equiv 0 \pmod{p}$, (III) is linear in y and, therefore, $b \equiv 0$ or $\not\equiv 0$.

Elimination of x' and z'' between (IV), (V), and (VI) gives

$$a'd^2 \equiv d'(ad - bc) \pmod{p}.$$

Since z'' is prime to p , (V) gives $d \not\equiv 0 \pmod{p}$, so that $ad - bc \not\equiv 0 \pmod{p}$. We may place $b = 0$, $\alpha = p$, $\beta = 0$, $c = 0$, $d = 1$, then a will take the values $1, 2, 3, \dots, p-1$ giving $p-1$ types.

D_1 .

The congruences for this case are

$$d'(yz' - y'z) \equiv \beta y' \pmod{p}, \quad (\text{I})$$

$$\alpha_1 x + \beta x' \equiv 0 \pmod{p}, \quad (\text{II})$$

$$d'yz'' \equiv by' \pmod{p}, \quad (\text{III})$$

$$ax + bx' \equiv 0 \pmod{p}, \quad (\text{IV})$$

$$d'z'' \equiv d \pmod{p}, \quad (\text{V})$$

$$cx + dx' \equiv 0 \pmod{p}. \quad (\text{VI})$$

z is contained in (I) alone, and therefore $\beta \equiv 0$ or $\not\equiv 0 \pmod{p}$.

(III) is linear in y , and $b \equiv 0$ or $\not\equiv 0 \pmod{p}$.

(V) gives $d \not\equiv 0 \pmod{p}$.

Elimination of x' between (II) and (VI) gives $\alpha_1 d - \beta c \equiv 0 \pmod{p}$, and between (IV) and (VI) gives $ad - bc \equiv 0 \pmod{p}$. The type group is derived by placing $a = 0$, $b = 0$, $\alpha = 0$, $\beta = 0$, $c = 0$ and $d = 1$.

Section 2.

1. *Groups with dependent generators.* In this section, G is generated by Q_1 and P where

$$Q_1^{p^2} = P^{hp^2}. \quad (1)$$

There is in G , a subgroup H_1 , of order p^{m-2} , which contains $\{P\}$ self-conjugately.²⁰ H_1 either contains, or does not contain Q_1^p . We will consider the second possibility in the present section, reserving the first for the next section.

2. *Determination of H_1 .* H_1 is generated by P and some other operator R_1 of G . R_1^p is contained in $\{P\}$. Let

$$R_1^p = P^{lp}. \quad (2)$$

Since $\{P\}$ is self-conjugate in H_1 ,²¹

$$R_1^{-1} P R_1 = P^{1+kp^{m-4}} \quad (3)$$

Denoting $R_1^a P^b R_1^c P^d \dots$ by the symbol $[a, b, c, d, \dots]$ we derive from (3)

$$[-y, x, y] = [0, x(1 + kyp^{m-4})] \quad (m > 4), \quad (4)$$

and

$$[y, x]^s = \left[sy, x \left\{ s + k \binom{s}{2} yp^{m-4} \right\} \right] \quad (5)$$

Placing $s = p$ and $y = 1$ in (5) we have, from (2)

$$[R_1 P^x]^p = R_1^p P^{xp} = P^{(l+x)p}.$$

Choosing x so that

$$x + l \equiv 0 \pmod{p^{m-4}},$$

$R = R_1 P^x$ is an operator of order p , which will be used in the place of R_1 , and $H = \{R, P\}$ with $R^p = 1$.

3. *Determination of H_2 .* We will now use the symbol $[a, b, c, d, e, f, \dots]$ to denote $Q_1^a R^b P^c Q_1^d R^e P^f \dots$.

H_1 and Q_1 generate G and all the operations of G are given by $[x, y, z]$ ($z = 0, 1, 2, \dots, p^2 - 1$; $y = 0, 1, 2, \dots, p - 1$; $x = 0, 1, 2, \dots, p^{m-3} - 1$), since these are p^m in number and are all distinct. There is in G a subgroup H_2 of order p^{m-1} which contains H_1 self-conjugately. H_2 is generated by H_1 and

²⁰BURNSIDE, *Theory of Groups*, Art. 54, p. 64.

²¹BURNSIDE, *Theory of Groups*, Art. 56, p. 66.

some operator $[z, y, x]$ of G . Q_1^z is then in H_2 and H_2 is the subgroup $\{Q_1^p, H_1\}$. Hence,

$$Q_1^{-p} P Q_1^p = R^\beta P^{\alpha_1}, \quad (6)$$

$$Q_1^{-p} P Q_1^p = R^{b_1} P^{ap^{m-4}}. \quad (7)$$

To determine α_1 and β we find from (6), (5) and (7)

$$\begin{aligned} [-p^2, 0, 1, p^2] = & \left[0, \frac{\alpha_1^p - b_1^p}{\alpha_1 - b_1} \beta, \alpha_1^p \left\{ 1 + \frac{\beta k}{2} \frac{\alpha_1^p - 1}{\alpha_1 - 1} p^{m-4} \right\} \right. \\ & \left. + a\beta \left\{ p \frac{\alpha_1^{p-1}}{\alpha_1 - b_1} - \frac{\alpha_1^p - b_1^p}{(\alpha_1 - b_1)^2} \right\} p^{m-4} \right]. \end{aligned}$$

By (1)

$$Q_1^{-p^2} P Q_1^{p^2} = P,$$

and, therefore,

$$\begin{aligned} \frac{\alpha_1^p - b_1^p}{\alpha_1 - b_1} \beta &\equiv 0 \pmod{p}, \\ \alpha_1^p &\equiv 1 \pmod{p^{m-4}}, \quad \text{and} \quad \alpha_1 \equiv 1 \pmod{p^{m-5}} \quad (m > 5). \end{aligned}$$

Let $\alpha_1 = 1 + \alpha_2 p^{m-5}$ and equation (6) is replaced by

$$Q_1^{-p} P Q_1^p = R^\beta P^{1+\alpha_2 p^{m-5}}. \quad (8)$$

To find a and b_1 we obtain from (7), (8) and (5)

$$[-p^2, 1, 0, p^2] = \left[0, b_1^p, a \frac{b_1^p - 1}{b_1 - 1} p^{m-4} \right].$$

By (1) and (4)

$$Q_1^{-p^2} R Q_1^{p^2} = P^{-lp^2} R P^{lp^2} = R,$$

and, hence,

$$b_1^p \equiv 1 \pmod{p}, \quad a \frac{b_1^p - 1}{b_1 - 1} \equiv 0 \pmod{p},$$

therefore $b_1 = 1$.

Substituting $b_1 = 1$ and $\alpha_1 = 1 + \alpha_2 p^{m-5}$ in the congruence determining α_1 we obtain $(1 + \alpha_2 p^{m-5})^p \equiv 1 \pmod{p^{m-3}}$, which gives $\alpha_2 \equiv 0 \pmod{p}$.

Let $\alpha_2 = \alpha p$ and equations (8) and (7) are now replaced by

$$Q_1^p P Q_1^p = R^\beta P^{1+\alpha p^{m-4}}, \quad (9)$$

$$Q_1^{-p} R Q_1^p = R P^{ap^{m-4}}. \quad (10)$$

From these we derive

$$[-yp, 0, x, yp] = \left[0, \beta xy, x + \{\alpha xy + a\beta x \binom{y}{2} + \beta k \binom{x}{2} y\} p^{m-4}\right], \quad (11)$$

$$[-yp, x, 0, yp] = [0, x, axyp^{m-4}]. \quad (12)$$

A continued use of (4), (11), and (12) yields

$$[zp, y, x]^s = [szp, sy + U_s, sx + V_s p^{m-4}] \quad (13)$$

where

$$\begin{aligned} U_s &= \beta \binom{s}{2} xz, \\ V_s &= \binom{s}{2} \left\{ \alpha xz + \beta k \binom{s}{2} z + kxy + ayz \right\} + \beta k \binom{s}{3} x^2 z \\ &\quad + \frac{1}{2} a\beta \left\{ \frac{1}{3!} s(s-1)(2s-1)z^2 - \binom{s}{2} z \right\}. \end{aligned}$$

4. Determination of G .

Since H_2 is self-conjugate in G_1 we have

$$Q_1^{-1} P Q_1 = Q_1^{\gamma p} R^\delta P^{\epsilon_1}, \quad (14)$$

$$Q_1^{-1} R Q_1 = Q_1^{cp} R^d P^{\epsilon p^{m-4}}. \quad (15)$$

From (14), (15) and (13)

$$[-p, 0, 1, p] = [\lambda p, \mu, \epsilon_1^p + \nu p^{m-4}]$$

and by (9) and (1)

$$\lambda p \equiv 0 \pmod{p^2}, \quad \epsilon_1^p + \nu p^{m-4} + \lambda h p \equiv 1 + \alpha p^{m-4} \pmod{p^{m-3}},$$

from which

$$\epsilon_1^p \equiv 1 \pmod{p^2}, \quad \text{and} \quad \epsilon_1 \equiv 1 \pmod{p} \quad (m > 5).$$

Let $\epsilon_1 = 1 + \epsilon_2 p$ and equation (14) is replaced by

$$Q_1^{-1} P Q_1 = Q_1^{\gamma p} R^\delta P^{1+\epsilon_2 p}. \quad (16)$$

From (15), (16), and (13)

$$[-p, 1, 0, p] = \left[c \frac{d^p - 1}{d - 1} p, d^p, K p^{m-4} \right]$$

where

$$K = \frac{d^p - 1}{d - 1} e + \sum_1^{p-1} a c d \frac{d^n (d^n - 1)}{2}.$$

By (10)

$$d^p \equiv 1 \pmod{p}, \quad \text{and} \quad d = 1$$

and by (1)

$$chp^2 \equiv ap^{m-4} \pmod{p^{m-3}}.$$

Equation (15) is now replaced by

$$Q_1^{-1} R Q_1 = Q_1^{cp} R P^{\epsilon p^{m-4}}. \quad (17)$$

A combination of (17), (16) and (13) gives

$$[-p, 0, 1, p] = \left[\left\{ \gamma \frac{(1 + \epsilon_2 p)^p - 1}{\epsilon_2 p^2} + c\delta \frac{p-1}{2} \right\} p^2, 0, (1 + \epsilon_2 p)^p \right].$$

By (9)

$$\left\{ \gamma \frac{(1 + \epsilon_2 p)^p - 1}{\epsilon_2 p^2} + c\delta \frac{p-1}{2} \right\} h p^2 + (1 + \epsilon_2 p)^p \equiv 1 + \alpha p^{m-4} \pmod{p^{m-3}},$$

$$\beta \equiv 0 \pmod{p}.$$

A reduction of the first congruence gives

$$\frac{(1 + \epsilon_2 p)^p - 1}{\epsilon p^2} \{ \epsilon_2 + \gamma h \} p^2 \equiv \left\{ \alpha - a\delta \frac{p-1}{2} \right\} p^{m-4} \pmod{p^{m-3}}$$

and, since

$$\frac{(1 + \epsilon_2 p)^p - 1}{\epsilon_2 p^2} \equiv 1 \pmod{p}, \quad (\epsilon_2 + \gamma h) p^2 \equiv 0 \pmod{p^{m-4}}$$

and

$$(\epsilon_2 + \gamma h) p^2 \equiv \left(\alpha + \frac{a\delta}{2} \right) p^{m-4} \pmod{p^{m-3}}. \quad (18)$$

From (17), (16), (13) and (18)

$$[-y, x, 0, y] = \left[cxy p, x, \{ exy + ac \binom{x}{2} y \} p^{m-4} \right], \quad (19)$$

$$[-y, 0, x, y] = \left[x \{ \gamma y + c\delta \binom{y}{2} \} p, \delta xy, x(1 + \epsilon_2 y p) + \theta p^{m-4} \right] \quad (20)$$

where

$$\begin{aligned} \theta = & \left\{ e\delta x + a\delta\gamma x + \epsilon_2 \left(\alpha + \frac{a\delta}{2} \right) x \right\} \binom{y}{2} \\ & + \frac{1}{2} ac \left\{ \frac{1}{3!} y(y-1)(2y-1)\delta^2 - \binom{y}{2} \delta \right\} \\ & + \left\{ \alpha\gamma y + \delta ky + a\delta xy^2 + (ac\delta^2 y + ac\delta) \binom{y}{2} \right\} \binom{x}{2}. \end{aligned}$$

From (19), (20), (4) and (18)

$$\{Q_1 P^x\}^{p^2} = Q_1^{p^2} P^{xp^2} = P^{(h+x)p^2}.$$

If x be so chosen that

$$h + x \equiv 0 \pmod{p^{m-5}}$$

$Q = Q_1 P^x$ is an operator of order p^2 which will be used in place Q_1 and $Q^{p^2} = 1$.

Placing $h = 0$ in (18) we get

$$\epsilon_2 p^2 \equiv 0 \pmod{p^{m-4}}.$$

Let $\epsilon_2 = \epsilon p^{m-6}$ and equation (16) is replaced by

$$Q^{-1} P Q = Q^{\gamma p} R^{\delta} P^{1+\epsilon p^{m-5}} \quad (21)$$

The congruence

$$ap^{m-4} \equiv cp^2 \pmod{p^{m-3}}$$

becomes

$$ap^{m-4} \equiv 0 \pmod{p^{m-3}}, \quad \text{and} \quad a \equiv 0 \pmod{p}.$$

Equations (19) and (20) are replaced by

$$[-y, x, 0, y] = [cxyp, x, exyp^{m-4}] \quad (22)$$

$$[-y, 0, x, y] = \left[\left\{ \gamma y + c\delta \binom{y}{2} \right\} xp, \delta xy, x(1 + \epsilon yp^{m-5}) + \theta p^{m-4} \right] \quad (23)$$

where

$$\theta = e\delta x \binom{y}{2} + \left\{ \alpha\gamma y + \delta ky + \alpha c\delta \binom{y}{2} \right\} \binom{x}{2}.$$

A formula for any power of an operation of G is derived from (4), (22) and (23)

$$[z, y, x]^s = [sz + U_s p, sy + V_s, sx + W_s p^{m-5}] \quad (24)$$

where

$$\begin{aligned} U_s &= \binom{s}{2} \{ \gamma xz + cyz \} + \frac{1}{2} c\delta x \left\{ \frac{1}{3!} s(s-1)(2s-1)z^2 - \binom{s}{2} z \right\}, \\ V_s &= \delta \binom{s}{2} xz, \\ W_s &= \binom{s}{2} \left\{ \epsilon xz + [(a\gamma + \delta k) \binom{x}{2} z + eyz + kxy] p \right\} \\ &\quad + \binom{s}{3} \{ \epsilon\gamma x + \epsilon y + \delta kx \} xzp + \frac{1}{2} c\delta \epsilon \left\{ \frac{1}{2} (s-1)z^2 - z \right\} \binom{s}{3} xp \\ &\quad + \frac{1}{2} \left\{ \delta ex + \alpha c\delta \binom{x}{2} \right\} \left\{ \frac{1}{3!} s(s-1)(2s-1)z^2 - \binom{s}{2} z \right\} p. \end{aligned}$$

5. *Transformations of the groups.* Placing $y = 1$ and $x = -1$ in (22) we obtain (17) in the form

$$R^{-1} Q R = Q^{1-cp} P^{-ep^{m-4}}.$$

A comparison of the generational equations of the present section with those of Section 1, shows that groups, in which $\delta \equiv 0 \pmod{p}$, are simply isomorphic with those in Section 1, so we need consider only those cases in which $\delta \not\equiv 0 \pmod{p}$.

All groups of this section are given by

$$G : \begin{cases} R^{-1} P R = P^{1+kp^{m-4}}, \\ Q^{-1} P Q = Q^{\gamma p} R^{\delta} P^{1+\epsilon p^{m-5}}, \\ Q^{-1} R Q = Q^{cp} R P^{\epsilon p^{m-4}}. \end{cases} \quad (25), (26), (27)$$

$R^p = 1$, $Q^{p^2} = 1$, and $P^{p^{m-3}} = 1$, ($k, \gamma, c, e = 0, 1, 2, \dots, p-1$; $\delta = 1, 2, \dots, p-1$; $\epsilon = 0, 1, 2, \dots, p^2-1$).

Not all these groups, however, are distinct. Suppose that G and G' of the above set are simply isomorphic and that the correspondence is given by

$$C = \begin{bmatrix} R, & Q, & P \\ R'_1, & Q'_1, & P'_1 \end{bmatrix}.$$

Since $R^p = 1$, $Q^{p^2} = 1$, and $P^{p^{m-3}} = 1$, $R'_1{}^p = 1$, $Q'_1{}^{p^2} = 1$ and $P'_1{}^{p^{m-3}} = 1$.

The forms of these operators are then

$$\begin{aligned} P'_1 &= Q'^z R'^y P'^x, \\ R'_1 &= Q'^{z'p} R'^{y'} P'^{x'p^{m-4}}, \\ Q'_1 &= Q'^{z''} R'^{y''} P'^{x''p^{m-5}}, \end{aligned}$$

where $dv[x, p] = 1$.

Since R is not contained in $\{P\}$, and Q^p is not contained in $\{R, P\}$ R'_1 is not contained in $\{P'_1\}$, and Q'^{p^2} is not contained in $\{R'_1, P'_1\}$.

Let

$$R'^{s'} = P'^{s'p^{m-4}}.$$

This becomes in terms of Q' , R' and P'

$$[s'z'p, s'y', s'x'p^{m-4}] = [0, 0, s'x'p^{m-4}],$$

and

$$s'y' \equiv 0 \pmod{p}, \quad s'z' \equiv 0 \pmod{p}.$$

Either y' or z' is prime to p or s' may be taken = 1.

Let

$$Q_1^{s''p} = R_1^{s'} P_1^{s'p^{m-4}},$$

and in terms of Q' , R' and P'

$$[s''z''p, 0, s''x''p^{m-4}] = [s'z'p, s'y', (s'x' + sx)p^{m-4}],$$

from which

$$s''z'' \equiv s'z' \pmod{p}, \quad \text{and} \quad s'y' \equiv 0 \pmod{p}.$$

Eliminating s' we find

$$s''y'z'' \equiv 0 \pmod{p},$$

$dv[y'z'', p] = 1$ or s'' may be taken = 1. We have then z'' , y' and x prime to p .

Since R , Q and P satisfy equations (25), (26) and (27) R_1 , Q_1 and P_1 do also. These become in terms of R' , Q' and P' .

$$\begin{aligned} [z + \Phi'_1 p, y, x + \Theta'_1 p^{m-4}] &= [z, y, x(1 + kp^{m-4})], \\ [z + \Phi'_2 p, y + \delta'xz'', x + \Theta'_2 p^{m-5}] &= [z + \Phi_2 p, y + \delta y', x + \Theta_2 p^{m-5}], \\ [(z' + \Phi'_3)p, y', \Theta'_3 p^{m-4}] &= [(z' + \Phi_3)p, y, \Theta_3 p^{m-4}], \end{aligned}$$

where

$$\begin{aligned} \Phi'_1 &= -c'yz', \quad \Theta'_1 = \epsilon'xz' + k'xy' - e'y'z, \\ \Phi'_2 &= \left\{ \gamma'z'' + c'\delta'\binom{z}{2} \right\} x + c'(yz'' - y''z), \\ \Theta'_2 &= \epsilon'xz'' + \left\{ \binom{x}{2} [\alpha'\gamma'z'' + \alpha'c'\delta'\binom{z''}{2}] + \delta'k'z'' \right. \\ &\quad \left. + \delta'e'x\binom{z''}{2} + e'(yz'' - y''z) + k'xy'' \right\} p, \\ \Phi_2 &= \gamma z'' + \delta z' + c'\delta y'z, \quad \Theta_2 \equiv \epsilon x + (\gamma x'' + \delta x + e'\delta y'z)p, \\ \Phi'_3 &= c'y'z'', \quad \Theta'_3 = e'y'z'', \quad \Phi_3 = cz'', \quad \Theta_3 = \epsilon x + cx''. \end{aligned}$$

A comparison of the members of these equations give seven congruences

$$\begin{aligned} \Phi'_1 &\equiv 0 \pmod{p}, & \text{(I)} \\ \Theta'_1 &\equiv kx \pmod{p}, & \text{(II)} \\ \Phi'_2 &\equiv \Phi_2 \pmod{p}, & \text{(III)} \\ \delta'xz'' &\equiv \delta y' \pmod{p}, & \text{(IV)} \\ \Theta'_2 &\equiv \Theta_2 \pmod{p^2}, & \text{(V)} \\ \Phi'_3 &\equiv cz'' \pmod{p}, & \text{(VI)} \\ \Theta'_3 &\equiv \Theta_3 \pmod{p}. & \text{(VII)} \end{aligned}$$

The necessary and sufficient condition for the simple isomorphism of G and G' is, that these congruences be consistent and admit of solution for the nine indeterminants with x, y' , and z'' prime to p .

Let $\kappa = \kappa_1 p^{\kappa_2}$, $dv[\kappa_1, p] = 1$ ($\kappa = k, \delta, \gamma, \epsilon, c, e$).

The groups are divided into three parts and each part is subdivided into 16 cases.

The methods used in discussing the congruences are the same as those used in Section 1.

6. *Reduction to types.* The three parts are given by

I.

	ϵ_2	δ_2
A	0	0
B	1	0
C	2	0

The subdivision into cases and the results of the discussion of the congruences are given in Table II.

II.

	k_2	γ_2	c_2	e_2	A	B	C
1	1	1	1	1			B_1
2	0	1	1	1			B_2
3	1	0	1	1	A_2	B_1	B_1
4	1	1	0	1			B_4
5	1	1	1	0			B_5
6	0	0	1	1	*	B_2	B_2
7	0	1	0	1	A_4		B_7
8	0	1	1	0	A_5	B_5	B_5
9	1	0	0	1	A_4	B_4	B_4
10	1	0	1	0	A_5	B_5	B_5
11	1	1	0	0	A_4	B_4	B_4
12	0	0	0	1	A_4	B_7	B_7
13	0	0	1	0	A_5	B_5	B_5
14	0	1	0	0	A_4	B_7	B_7
15	1	0	0	0	A_4	B_4	B_4
16	0	0	0	0	A_4	B_7	B_7

A_6 divides into two parts.

The groups of A_6 in which $\delta k + \epsilon \gamma \equiv 0 \pmod{p}$ are simply isomorphic with the groups of A_1 and those in which $\delta k + \epsilon \gamma \not\equiv 0 \pmod{p}$ are simply isomorphic with the groups of A_2 . The types are given by equations (25), (26) and (27) where the constants have the values given in Table III.

III.

	k	δ	γ	ϵ	c	e
A_1	0	1	0	1	0	0
A_2	1	1	0	1	0	0
A_4	0	1	0	1	1	0
A_5	0	1	0	1	0	ω
B_1	0	1	0	p	0	0
B_2	1	1	0	p	0	0
B_4	0	1	0	p	1	0
B_5	0	1	0	p	0	κ
B_7	1	1	0	p	ω	0

$\kappa = 1$, and a non-residue \pmod{p} ,
 $\omega = 1, 2, \dots, p-1$.

A detailed analysis of several cases is given below, as a general illustration of the methods used.

A_1 .

The special forms of the congruences for this case are

$$\epsilon'xz' \equiv kx \pmod{p}, \tag{II}$$

$$\gamma z'' + \delta z' \equiv 0 \pmod{p}, \tag{III}$$

$$\delta'xz'' \equiv \delta y' \pmod{p}, \tag{IV}$$

$$\epsilon'xz'' \equiv \epsilon x \pmod{p}, \tag{V}$$

$$cz'' \equiv 0 \pmod{p}, \tag{VI}$$

$$ex \equiv 0 \pmod{p}. \tag{VII}$$

Congruence (IV) gives $\delta \not\equiv 0 \pmod{p}$, from (II) k can be $\equiv 0$ or $\not\equiv 0 \pmod{p}$, (III) gives $\gamma \equiv 0$ or $\not\equiv 0$, (V) gives $\epsilon \not\equiv 0$, (VI) and (VII) give $c \equiv e \equiv 0 \pmod{p}$. Elimination of x , z' and z'' between (II), (III) and (V) gives $\delta k + \gamma \epsilon \equiv 0 \pmod{p}$. If $k \equiv 0$, then $\gamma \equiv 0 \pmod{p}$ and if $k \not\equiv 0$, then $\gamma \not\equiv 0 \pmod{p}$.

A_2 .

The congruences for this case are

$$\epsilon'xz' + k'xy' \equiv kx \pmod{p}, \tag{II}$$

$$\gamma x'' + \delta z' \equiv 0 \pmod{p}, \tag{III}$$

$$\delta'xz'' \equiv \delta y' \pmod{p}, \tag{IV}$$

$$\epsilon'xz'' \equiv \epsilon x \pmod{p}, \tag{V}$$

$$cz'' \equiv 0 \pmod{p}, \tag{VI}$$

$$ex \equiv 0 \pmod{p}. \tag{VII}$$

Congruence (III) gives $\gamma \equiv 0$ or $\not\equiv 0$, (IV) gives $\delta \not\equiv 0$, (V) $\epsilon \not\equiv 0$, (VI) and (VII) give $c \equiv e \equiv 0 \pmod{p}$. Elimination of x , z' , and z'' between (II), (III) and (V) gives

$$\delta k + \gamma \epsilon \equiv k' \delta y' \pmod{p}$$

from which

$$\delta k + \gamma \epsilon \not\equiv 0 \pmod{p}.$$

If $k \equiv 0$, then $\gamma \not\equiv 0$, and if $\gamma \equiv 0$ then $k \not\equiv 0 \pmod{p}$.

Both γ and k can be $\not\equiv 0 \pmod{p}$ provided the above condition is fulfilled.

A_5 .

The congruences for this case are

$$\epsilon' x z' - e' y' z \equiv k x \pmod{p}, \quad (\text{II})$$

$$\gamma z'' + \delta z' \equiv 0 \pmod{p}, \quad (\text{III})$$

$$\delta' x z'' \equiv \delta y' \pmod{p}, \quad (\text{IV})$$

$$\epsilon' x z'' \equiv e x \pmod{p}, \quad (\text{V})$$

$$c z'' \equiv 0 \pmod{p}, \quad (\text{VI})$$

$$e' y' z'' \equiv e x \pmod{p}. \quad (\text{VII})$$

(II) and (III) are linear in z and z' so k and γ are \equiv or $\not\equiv 0 \pmod{p}$ independently, (IV) gives $\delta \not\equiv 0$, (V) give $\epsilon \not\equiv 0$, (VI) $c \equiv 0$, and (VII) $e \not\equiv 0$.

Elimination between (IV), (V), and (VII) gives

$$\delta' e' \epsilon^2 \equiv \delta e \epsilon'^2 \pmod{p}.$$

The types are derived by placing $\epsilon = \delta = 1$, and $e = 1, 2, \dots, p-1$.

B_5 .

The congruences for this case are

$$-e' y' z \equiv k x \pmod{p}, \quad (\text{II})$$

$$\gamma z'' + \delta z' \equiv 0 \pmod{p}, \quad (\text{III})$$

$$\delta' x z'' \equiv \delta y' \pmod{p}, \quad (\text{IV})$$

$$\epsilon'_1 x z'' + \delta' e' x \binom{z''}{2} + e' y z'' \equiv e_1 x + \gamma x'' + \delta x' \pmod{p}, \quad (\text{V})$$

$$c z'' \equiv 0 \pmod{p}, \quad (\text{VI})$$

$$e' y' z'' \equiv e x \pmod{p}. \quad (\text{VII})$$

(II), and (III) being linear in z and z' give $k \equiv 0$ or $\not\equiv 0$, and $\gamma \equiv 0$ or $\not\equiv 0 \pmod{p}$, (IV) gives $\delta \not\equiv 0$, (V) being linear in x' gives $\epsilon_1 \equiv 0$ or $\not\equiv 0 \pmod{p}$, (VI) gives $c \equiv 0$ and (VII) $e \not\equiv 0$.

Elimination of x and y' from (IV) and (VII) gives

$$\delta' e' z''^2 \equiv \delta e \pmod{p}.$$

δe is a quadratic residue or non-residue \pmod{p} according as $\delta' e'$ is a residue or non-residue.

The two types are given by placing $\delta = 1$, and $e = 1$ and a non-residue \pmod{p} .

Section 3.

1. *Groups with dependent generators continued.* As in Section 2, G is here generated by Q_1 and P , where

$$Q_1^{p^2} = P^{hp^2}.$$

Q_1^p is contained in the subgroup H_1 of order p^{m-2} , $H_1 = \{Q_1^p, P\}$.

2. *Determination of H_1 .* Since $\{P\}$ is self-conjugate in H_1

$$Q_1^{-p} P Q_1^p = P^{1+kp^{m-4}}. \quad (1)$$

Denoting $Q_1^a P^b Q_1^c P^d \dots$ by the symbol $[a, b, c, d, \dots]$, we have from (1)

$$[-yp, x, yp] = [0, x(1+kyp^{m-4})] \quad (m > 4). \quad (2)$$

Repeated multiplication with (2) gives

$$[yp, x]^s = \left[syp, x \left\{ s + k \binom{s}{2} yp^{m-4} \right\} \right]. \quad (3)$$

3. *Determination of H_2 .* There is a subgroup H_2 of order p^{m-1} which contains H_1 self-conjugately.²² H_2 is generated by H_1 and some operator R_1 of G . R_1^p is contained in H_1 , in fact in $\{P\}$, since if $R_1^{p^2}$ is the first power of R_1 in $\{P\}$, then $H_2 = \{R_1, P\}$, which case was treated in Section 1.

$$R_1^p = P^{lp}. \quad (4)$$

Since H_1 is self-conjugate in H_2

$$R_1^{-1} P R_1 = Q_1^{\beta p} P^{\alpha_1}, \quad (5)$$

$$R_1^{-1} Q^p R_1 = Q_1^{bp} P^{\alpha_1 p}. \quad (6)$$

Using the symbol $[a, b, c, d, e, f, \dots]$ to denote $R_1^a Q_1^b P^c R_1^d Q_1^e P^f \dots$, we have from (5), (6) and (3)

$$[-p, 0, 1, p] = [0, \beta Np, \alpha_1^p + Mp], \quad (7)$$

²²BURNSIDE, *Theory of Groups*, Art. 54, p. 64.

and by (4)

$$\alpha_1^p \equiv 1 \pmod{p}, \quad \text{and} \quad \alpha_1 \equiv 1 \pmod{p}.$$

Let $\alpha_1 = 1 + \alpha_2 p$ and (5) is now replaced by

$$R_1^{-1} P R_1 = Q_1^{\beta p} P^{1+\alpha_2 p}. \quad (8)$$

From (6), (8) and (3)

$$[-p, p, 0, p] = [0, b^p p, a_1 \frac{b^p - 1}{b - 1} p + a_1 U p^2],$$

and by (4) and (2)

$$R_1^{-p} Q_1^p R_1^p = Q_1^p$$

and therefore $b^p \equiv 1 \pmod{p}$, and $b = 1$. Placing $b = 1$ in the above equation the exponent of P takes the form

$$a_1 p^2 (1 + U' p) = a_1 \frac{\{1 + (\alpha_2 + \beta h)p\}^p - 1}{(\alpha_2 + \beta h)p^2} p^2$$

from which

$$a_1 p^2 (1 + U' p) \equiv 0 \pmod{p^{m-3}}$$

or

$$a_1 \equiv 0 \pmod{p^{m-5}} \quad (m > 5).$$

Let $a_1 = a p^{m-5}$ and (6) is replaced by

$$R_1^{-1} Q_1^p R_1 = Q_1^p P^{a p^{m-4}}. \quad (9)$$

(7) now has the form

$$[-p, 0, 1, p] = [0, \beta N p, (1 + \alpha_2 p)^p + M p^2],$$

where

$$N = p \quad \text{and} \quad M = \beta h \left\{ \frac{(1 + \alpha_2 p)^p - 1}{\alpha_2 p^2} - 1 \right\},$$

from which

$$(1 + \alpha_2 p)^p + \frac{(1 + \alpha_2 p)^p - 1}{\alpha_2 p^2} \beta h p^2 \equiv 1 \pmod{p^{m-3}}$$

or

$$\frac{(1 + \alpha_2 p)^p - 1}{\alpha_2 p^2} \{\alpha_2 + \beta h\} p^2 \equiv 0 \pmod{p^{m-3}}$$

and since

$$\begin{aligned} \frac{(1 + \alpha_2 p)^p - 1}{\alpha_2 p^2} &\equiv 1 \pmod{p} \\ (\alpha_2 + \beta h) p^2 &\equiv 0 \pmod{p^{m-3}}. \end{aligned} \quad (10)$$

From (8), (9), (10) and (3)

$$[-y, 0, x, y] = [0, \beta x y p, x(1 + \alpha_2 y p) + \theta p^{m-4}], \quad (11)$$

$$[-y, x p, 0, y] = [0, x p, a x y p^{m-4}], \quad (12)$$

where

$$\theta = a \beta x \binom{y}{2} + \beta k \binom{x}{2} y.$$

By continued use of (11), (12), (2) and (10)

$$[z, y p, x]^s = [s z, (s y + U_s) p, x s + V_s p], \quad (13)$$

where

$$\begin{aligned} U_s &= \beta \binom{s}{2} x z \\ V_s &= \binom{s}{2} \left\{ \alpha_2 x z + [a y z + k x y + \beta k \binom{x}{2} z] p^{m-5} \right\} \\ &\quad + \left\{ \beta \binom{s}{3} x^2 z + \frac{1}{2} a \beta \left[\frac{1}{3!} s(s-1)(2s-1) z^2 - \binom{s}{2} z \right] x \right\} p^{m-5}. \end{aligned}$$

Placing in this $y = 0$, $z = 1$ and $s = p$,²³

$$(R_1 P^x)^p = R_1^p P^{x p} = P^{(x+l)p},$$

determine x so that

$$x + l \equiv 0 \pmod{p^{m-4}},$$

then $R = R_1 P^x$ is an operator of order p which will be used in the place of R_1 , $R^p = 1$.

4. *Determination of G .* Since H_2 is self-conjugate in G

$$Q_1^{-1} P Q_1 = R^\gamma Q_1^{\delta p} P^{\epsilon_1}, \quad (14)$$

$$Q_1^{-1} R Q_1 = R^c Q_1^{d p} P^{\epsilon_1 p}. \quad (15)$$

²³Terms of the form $(Ax^2 + Bx)p^{m-4}$ in the exponent of P for $p = 3$ and $m > 5$ do not alter the result.

From (15)

$$(R^c Q_1^{dp} P^{e_1 p})^p = 1,$$

by (13)

$$Q_1^{dp^2} P^{e_1 p^2} = P^{(e_1 + dh)p^2} = 1,$$

and

$$(e_1 + dh)p^2 \equiv 0 \pmod{p^{m-3}}. \quad (16)$$

From (14), (15) and (13)

$$[0, -p, 1, 0, p] = [L, Mp, \epsilon_1^p + Np]. \quad (17)$$

By (1)

$$\epsilon_1^p \equiv 1 \pmod{p}, \quad \text{and} \quad \epsilon_1 \equiv 1 \pmod{p}.$$

Let $\epsilon_1 = 1 + \epsilon_2 p$ and (14) is replaced by

$$Q_1^{-1} P Q_1 = R^\gamma Q_1^{\delta p} P^{1 + \epsilon_2 p}. \quad (18)$$

From (15), (18), and (13)

$$[0, -p, 0, 1, p] = \left[c_p, \frac{c^p - 1}{c - 1} dp, Kp \right].$$

Placing $x = 1$ and $y = -1$ in (12) we have

$$[0, -p, 0, 1, p] = [1, 0, -ap^{m-4}], \quad (19)$$

and therefore $c^p \equiv 1 \pmod{p}$, and $c = 1$. (15) is now replaced by

$$Q_1^{-1} R Q_1 = R Q_1^{dp} P^{e_1 p}. \quad (20)$$

Substituting $1 + \epsilon_2 p$ for ϵ_1 and 1 for c in (17) gives, by (16)

$$[0, -p, 1, p] = [0, Mp^2, (1 + \epsilon_2 p)^p + Np^2],$$

where

$$M = \gamma d \frac{p-1}{2} + \delta \frac{(1 + \epsilon_2 p)^p - 1}{\epsilon_2 p^2}$$

and

$$N = \frac{e_1 \gamma}{(\epsilon_2 + \delta h)p^2} \left\{ \frac{[1 + (\epsilon_2 + \delta h)p]^p - 1}{(\epsilon_2 + \delta h)p} - p \right\}.$$

By (1)

$$(1 + \epsilon_2 p)^p + (N + Mh)p^2 \equiv 1 + kp^{m-4} \pmod{p^{m-3}},$$

or reducing

$$\psi(\epsilon_2 + \delta h)p^2 \equiv kp^{m-4} \pmod{p^{m-3}},$$

where

$$\psi = \frac{(1 + \epsilon_2 p)^p - 1}{\epsilon_2 p^2} + N - e_1 \gamma \frac{p-1}{2}.$$

Since

$$\begin{aligned} \psi &\equiv 1 \pmod{p}. \\ (\epsilon_2 + \delta h)p^2 &\equiv kp^{m-4} \pmod{p^{m-3}}. \end{aligned} \quad (21)$$

From (18), (20), (13), (16) and (21)

$$[0, -y, x, 0, y] = [\gamma xy, \theta_1 p, x + \phi_1 p], \quad (22)$$

$$[0, -y, 0, x, y] = [x, dxy p, \phi_2 p], \quad (23)$$

where

$$\begin{aligned} \theta_1 &= d\gamma x \binom{y}{2} + \delta xy + \beta \gamma \binom{x}{2} y, \\ \phi_1 &= \epsilon_2 xy + \alpha_2 \gamma \binom{x}{2} y + e_1 \gamma \binom{y}{2} x + \left\{ x \binom{y}{2} (\epsilon_2 k + \delta \gamma) \right. \\ &\quad \left. + \frac{1}{2} ad \left[\frac{1}{3!} y(y-1)(2y-1)\gamma^2 - \frac{y}{2} \gamma \right] x + a\gamma^2 dx \frac{1}{3!} y(y+1)(y-1) \right. \\ &\quad \left. + e_1 \gamma k \binom{y}{3} x + \frac{1}{2} a\beta \left[\frac{1}{3!} x(x-1)(2x-1)\gamma^2 y^2 - \binom{x}{2} \gamma y \right] \right. \\ &\quad \left. + \binom{x}{2} (a+k) [dy \binom{y}{2} + \delta y] + \beta \gamma \binom{x}{3} \right\} p^{m-5}, \\ \phi_2 &= e_1 xy + \left\{ e_1 k \binom{y}{2} + ad \binom{x}{2} y \right\} p^{m-5}. \end{aligned}$$

Placing $x = 1$ and $y = p$ in (23) and by (16)

$$Q_1^{-p} R Q_1^p = R,$$

and by (19)

$$a \equiv 0 \pmod{p}.$$

A continued multiplication, with (11), (22), and (23), gives

$$(Q_1 P^x)^{p^2} = Q_1^{p^2} P^{xp^2} = P^{(x+l)p^2}.$$

Let x be so chosen that

$$(x + l) \equiv 0 \pmod{p^{m-5}},$$

then $Q = Q_1 P^x$ is an operator of order p^2 which will be used in place of Q_1 , $Q^{p^2} = 1$ and

$$h \equiv 0 \pmod{p^{m-5}}.$$

From (21), (10) and (16)

$$\epsilon_2 p^2 \equiv k p^{m-4}, \quad \alpha_2 p^2 \equiv 0 \quad \text{and} \quad e_1 p^2 \equiv 0 \pmod{p^{m-3}}.$$

Let $\epsilon_2 = \epsilon p^{m-6}$, $\alpha_2 = \alpha p^{m-5}$ and $e_1 = e p^{m-5}$. Then equations (18), (20) and (8) are replaced by

$$G : \begin{cases} Q^{-1} P Q = R^\gamma Q^{\delta p} P^{1+\epsilon p^{m-5}}, \\ Q^{-1} R Q = R Q^{\delta p} P^{\epsilon p^{m-4}}, \\ R^{-1} P R = Q^{\beta p} P^{1+\alpha p^{m-4}}, \\ R^p = 1, \quad Q^{p^2} = 1, \quad P^{p^{m-3}} = 1. \end{cases} \quad (24), (25), (26)$$

(11), (22) and (23) are replaced by

$$[-y, 0, x, y] = [0, \beta x y p, x + \phi p^{m-4}], \quad (27)$$

$$[0, -y, x, 0, y] = [\gamma x y, \theta_1 p, x + \phi_1 p^{m-5}], \quad (28)$$

$$[0, -y, 0, x, y] = [x, d x y p, \phi_2 p^{m-4}], \quad (29)$$

where

$$\begin{aligned} \phi &= \alpha x y + \beta k \binom{x}{2} y, \quad \theta_1 = d \gamma \binom{y}{2} x + \delta x y + \beta \gamma \binom{x}{2} y, \\ \phi_1 &= e x y + \left\{ e \gamma x \binom{y}{2} + \binom{x}{2} (\alpha \gamma y + d \gamma k \binom{y}{2}) + \delta k y \right\} p, \\ \phi_2 &= e x y. \end{aligned}$$

A formula for a general power of any operator of G is derived from (27), (28) and (29)

$$[0, z, 0, y, 0, z]^s = [0, s z + U_s p, 0, s y + V_s, 0, s x + W_s p^{m-5}], \quad (30)$$

where

$$\begin{aligned} U_s &= \binom{s}{2} \left\{ \delta x z + d y z + \beta x y + \beta \gamma \binom{x}{2} z \right\} \\ &\quad + \frac{1}{2} d x \left\{ \frac{1}{3!} s(s-1)(2s-1) z^2 - \binom{s}{2} z \right\} x + \beta \gamma \binom{s}{2} x^2 z, \\ V_s &= \gamma \binom{s}{2} x z, \end{aligned}$$

$$\begin{aligned}
W_s = & \binom{s}{2} \{ \epsilon xz + [axy + eyz + (\beta ky + \alpha\beta\gamma + \delta kz) \binom{x}{2}] p \} \\
& + \binom{s}{3} \{ \alpha\gamma x^2 z + dkxyz + \delta kx^2 z + \beta kx^2 y + 2\beta\gamma k \binom{x}{2} xz \} p \\
& + \beta\gamma k \binom{s}{4} x^3 z p + \frac{1}{2} \left\{ \frac{1}{3!} s(s-1)(2s-1)z^2 - \frac{s}{2}z \right\} \{ e\gamma x + d\gamma k \binom{x}{2} \} p \\
& + \frac{1}{2} d\gamma k \left[\frac{1}{2}(s-1)z^2 - z \right] \binom{s}{3} x^2.
\end{aligned}$$

A comparison of the generational equations of the present section with those of Sections 1 and 2, shows that, $\gamma \equiv 0 \pmod{p}$ gives groups simply isomorphic with those of Section 1, while $\beta \equiv 0 \pmod{p}$, groups simply isomorphic with those of Section 2 and we need consider only the groups in which β and γ are prime to p .

5. *Transformation of the groups.* All groups of this section are given by equations (24), (25), and (26), where $\gamma, \beta = 1, 2, \dots, p-1$; $\alpha, \delta, d, e = 0, 1, 2, \dots, p-1$; and $\epsilon = 0, 1, 2, \dots, p^2-1$.

Not all of these, however are distinct. Suppose that G is simply isomorphic with G' and that the correspondence is given by

$$C = \begin{bmatrix} R, & Q, & P \\ R'_1, & Q'_1, & P'_1 \end{bmatrix}.$$

An inspection of (30) gives

$$\begin{aligned}
R'_1 &= Q'^{z'p} R'^{y''} P'^{x''p^{m-4}}, \\
Q'_1 &= Q'^{z'} R'^{y'} P'^{x'p^{m-5}}, \\
P'_1 &= Q'^z R'^y P'^x,
\end{aligned}$$

with $dv[x, p] = 1$. Since Q^p is not in $\{P\}$, and R is not in $\{Q^p, P\}$, Q'^p_1 is not in $\{P'_1\}$ and R'_1 is not in $\{Q'^p_1, P'_1\}$. Let

$$Q'^{s'p}_1 = P'^{sp^{m-4}}_1.$$

This is in terms of R' , Q' , and P' ,

$$[0, s'z'p, s'x'p^{m-4}] = [0, 0, sxp^{m-4}].$$

From which

$$s'z'p \equiv 0 \pmod{p^2},$$

and z' must be prime to p , since otherwise s' can = 1. Let

$$R'^{s''}_1 = Q'^{s'p}_1 P'^{sp^{m-4}}_1,$$

or in terms of R' , Q' , and P' ,

$$[s''y'', s''z''p, s''x''p^{m-4}] = [0, s'z'p, (sx + s'x')p^{m-4}]$$

and

$$s''z'' \equiv s'z' \pmod{p}, \quad s''y'' \equiv 0 \pmod{p},$$

and y'' is prime to p , since otherwise s'' can = 1. Since R , Q , and P satisfy equations (24), (25) and (26), R'_1 , Q'_1 , and P'_1 must also satisfy them. These become when reduced in terms of R' , Q' and P'

$$\begin{aligned} & [0, z + \theta'_1 p, 0, y + \gamma'xz', 0, x + \psi'_1 p^{m-5}] \\ & \quad = [0, z + \theta_1 p, 0, y + \gamma y'', 0, x + \psi_1 p^{m-5}], \\ & [0, (z'' + \theta'_2)p, 0, y'', 0, (x'' + \psi_2)p^{m-4}] \\ & \quad = [0, (z'' + \theta_2)p, 0, y'', 0, (x'' + \psi_2)p^{m-4}], \\ & [0, z + \theta'_3 p, 0, y, 0, x + \psi'_3 p^{m-4}] = [0, z + \theta_3 p, 0, y, 0, x + \psi_3 p^{m-4}], \end{aligned}$$

where

$$\begin{aligned} \theta'_1 &= d'(yz' - y'z) + x \left\{ d'\gamma' \binom{z'}{2} + \delta'z' + \beta'y' \right\} + \beta'\gamma' \binom{x}{2} z', \\ \theta_1 &= \gamma z'' + \delta z' + d'\gamma y'' z, \\ \psi'_1 &= \epsilon'xz' + \left\{ e'\gamma'x \binom{z'}{2} + \binom{x}{2} \left[\alpha'\gamma'z' + \gamma'\epsilon'd'k' \binom{z'}{2} + \delta'\epsilon k'z' + \beta'k'y' \right] \right. \\ & \quad \left. + \beta'\gamma' \binom{x}{3} z' + e'(yz' - y'z) + \alpha'xy' \right\} p, \\ \psi_1 &= \epsilon x + \{ \delta x' + \gamma x'' + e'\gamma y'' z \} p, \\ \theta'_2 &= d'y''z', \quad \theta_2 = dz', \quad \psi'_2 = e'y''z, \quad \psi_2 = dx' + ex, \\ \theta'_3 &= \beta'xy'' - d'y''z, \quad \theta_3 = \beta z', \\ \psi_3 &= \epsilon'xz'' - e'y''z + \alpha'xy'' + \beta'\epsilon' \binom{x}{2} y'', \quad \psi_3 = \alpha x + \beta x'. \end{aligned}$$

A comparison of the two sides of these equations give seven congruences

$$\theta'_1 \equiv \theta_1 \pmod{p}, \tag{I}$$

$$\gamma'xz' \equiv \gamma y'' \pmod{p}, \tag{II}$$

$$\psi'_1 \equiv \psi_1 \pmod{p^2}, \tag{III}$$

$$\theta'_2 \equiv \theta_2 \pmod{p}, \tag{IV}$$

$$\psi'_2 \equiv \psi_2 \pmod{p}, \tag{V}$$

$$\theta'_3 \equiv \theta_3 \pmod{p}, \tag{VI}$$

$$\psi'_3 \equiv \psi_3 \pmod{p}. \tag{VII}$$

(VI) is linear in z provided $d' \not\equiv 0 \pmod{p}$ and z may be so determined that $\beta \equiv 0 \pmod{p}$ and therefore all groups in which $d' \not\equiv 0 \pmod{p}$ are simply isomorphic with groups in Section 2.

Consequently we need only consider groups in which $d' \equiv 0 \pmod{p}$.

As before we take for G' the simplest case and associate with it all simply isomorphic groups G . We then take as G' the simplest case left and proceed as above.

Let $\kappa = \kappa_1 p^{\kappa_2}$ where $dv[\kappa_1, p] = 1$, ($\kappa = \alpha, \beta, \gamma, \delta, \epsilon, d, e$).

For convenience the groups are divided into three sets and each set is subdivided into eight cases.

The sets are given by

$$\begin{aligned} A : \quad & \epsilon_2 = 0, \quad \beta_2 = 0, \quad \gamma_2 = 0, \\ B : \quad & \epsilon_2 = 1, \quad \beta_2 = 0, \quad \gamma_2 = 0, \\ C : \quad & \epsilon_2 = 2, \quad \beta_2 = 0, \quad \gamma_2 = 0. \end{aligned}$$

The subdivision into cases and results of the discussion are given in Table I.

I.

	δ_2	e_2	α_2	A	B	C
1	1	1	1			B_1
2	0	1	1	A_1	B_1	B_1
3	1	0	1			B_3
4	1	1	0	A_1	B_1	B_1
5	0	0	1	A_3	B_3	B_3
6	0	1	0	A_1	B_1	B_1
7	1	0	0	A_3	B_3	B_3
8	0	0	0	A_3	B_3	B_3

6. *Reduction to types.* The types of this section are given by equations (24), (25) and (26) with $\alpha = 0, \beta = 1, \lambda = 1$ or a quadratic non-residue \pmod{p} , $\delta \equiv 0; \epsilon = l, e = 0, 1, 2, \dots, p-1$; and $\epsilon = p, e = 0, 1$, or a non-residue \pmod{p} , $2p+6$ in all.

The special forms of the congruences for these cases are given below.

A_1 .

$$\beta' \gamma' \binom{x}{2} z' + \beta' xy' \equiv \gamma z'' + \delta z' \pmod{p}, \quad (\text{I})$$

$$\gamma' xz' \equiv \gamma y'' \pmod{p}, \quad (\text{II})$$

$$\epsilon' xz' \equiv \epsilon x \pmod{p}, \quad (\text{III})$$

$$dz' \equiv 0 \pmod{p}, \quad (\text{IV})$$

$$\epsilon x \equiv 0 \pmod{p}, \quad (\text{V})$$

$$\beta' xy'' \equiv \beta z' \pmod{p}, \quad (\text{VI})$$

$$\epsilon' xz'' + \beta' \epsilon' \binom{x}{2} y' \equiv \alpha x + \beta x' \pmod{p}. \quad (\text{VII})$$

(I) is linear in z'' and $\delta \equiv 0$ or $\not\equiv 0$, (II) gives $\gamma \not\equiv 0$, (III) $\epsilon \not\equiv 0$, (IV) and (V) $d \equiv e \equiv 0$, (VI) $\beta \not\equiv 0$, (VII) is linear in x' and $\alpha \equiv 0$ or $\not\equiv 0 \pmod{p}$.

Elimination of y'' and z' between (II) and (VI) gives

$$\beta'\gamma'x^2 \equiv \beta\gamma \pmod{p}$$

and $\beta\gamma$ is a residue or non-residue \pmod{p} according as $\beta'\gamma'$ is a residue or non-residue.

A_3 .

$$\beta'\gamma' \binom{x}{2} z' + \beta'xy' \equiv \gamma z'' + \delta z' \pmod{p}, \quad (\text{I})$$

$$\gamma'xz' \equiv \gamma y'' \pmod{p}, \quad (\text{II})$$

$$\epsilon'z' \equiv \epsilon \pmod{p}, \quad (\text{III})$$

$$d \equiv 0 \pmod{p}, \quad (\text{IV})$$

$$e'y''z' \equiv ex \pmod{p}, \quad (\text{V})$$

$$\beta'xy'' \equiv \beta z' \pmod{p}, \quad (\text{VI})$$

$$\epsilon'xz'' - e'y''z + \beta'\epsilon' \binom{x}{2} y' \equiv \alpha x + \beta x' \pmod{p}. \quad (\text{VII})$$

(I) is linear in z'' and $\delta \equiv 0$ or $\not\equiv 0$. (II) gives $\gamma \not\equiv 0$, (III) $\epsilon \not\equiv 0$, (V) $e \not\equiv 0$ and (VI) $\beta \not\equiv 0$. (VII) is linear in x' and $\alpha \equiv 0$ or $\not\equiv 0 \pmod{p}$.

Elimination between (II) and (VI) gives

$$\beta'\gamma'x^2 \equiv \beta\gamma \pmod{p},$$

and between (II), (III), and (IV) gives

$$\epsilon'^2\gamma e \equiv \epsilon^2\gamma'e' \pmod{p}.$$

$\beta\gamma$ is a residue, or non-residue, according as $\beta'\gamma'$ is or is not, and if γ and ϵ are fixed, e must take the $(p-1)$ values $1, 2, \dots, p-1$.

B_1 .

$$\beta'\gamma' \binom{x}{2} z' + \beta'xy' \equiv \gamma z'' + \delta z' \pmod{p}, \quad (\text{I})$$

$$\gamma'xz' \equiv \gamma y'' \pmod{p}, \quad (\text{II})$$

$$\epsilon'_1xz' + \beta'xz' \binom{x}{3} \equiv \epsilon_1x + \delta x' + \gamma x'' \pmod{p}, \quad (\text{III})$$

$$ex \equiv 0 \pmod{p}, \quad (\text{IV})$$

$$\beta'xy'' \equiv \beta z' \pmod{p}, \quad (\text{VI})$$

$$\alpha x + \beta x' \equiv 0 \pmod{p}. \quad (\text{VII})$$

(I) gives $\delta \equiv 0$ or $\not\equiv 0$, (II) $\gamma \not\equiv 0$, (III) is linear in x'' and gives $\epsilon_1 \equiv 0$ or $\not\equiv 0$, (V) $e = 0$, (VI) $\beta \not\equiv 0$ and (VII) is linear in x' and gives $\alpha \equiv 0$ or $\not\equiv 0$.

Elimination between (II) and (VI) gives

$$\beta'\gamma'x^2 \equiv \beta\gamma \pmod{p}.$$

B_3 .

$$\beta' \gamma' \binom{x}{2} z' + \beta' x y' \equiv \gamma \beta'' + \delta z' \pmod{p}, \quad (\text{I})$$

$$\gamma' x z' \equiv \gamma y' \pmod{p}, \quad (\text{II})$$

$$\begin{aligned} \epsilon'_1 x z' + e' \gamma' x \binom{z'}{2} + \beta' \gamma' \binom{x}{3} + e' (y z' - y' z) \\ \equiv \epsilon_1 x + \delta x' + \gamma x'' + e' \gamma z y'' \pmod{p}, \end{aligned} \quad (\text{III})$$

$$e' y'' z' \equiv e x \pmod{p}. \quad (\text{V})$$

$$\beta' x y'' \equiv \beta z' \pmod{p}, \quad (\text{VI})$$

$$-e' y'' z \equiv \alpha x + \beta x' \pmod{p}. \quad (\text{VII})$$

(I) gives $\delta \equiv 0$ or $\not\equiv 0$, (II) $\gamma \not\equiv 0$, (III) is linear in x'' and gives $\epsilon_1 \equiv 0$ or $\not\equiv 0$, (V) $e \not\equiv 0$, (VI) $\beta \not\equiv 0$, (VII) is linear in x' and gives $\alpha \equiv 0$ or $\not\equiv 0 \pmod{p}$. Elimination of y'' and z' between (II) and (VI) gives

$$\beta' \gamma' x^2 \equiv \beta \gamma \pmod{p},$$

and between (V) and (VI) gives

$$\beta' e' y''^2 \equiv \beta e \pmod{p}$$

and $\beta \gamma$ and βe are residues or non-residues, independently, according as $\beta' \gamma'$ and $\beta' e'$ are residues or non-residues.

Class III.

1. *General relations.* In this class, the p th power of every operator of G is contained in $\{P\}$. There is in G a subgroup H_1 of order p^{m-2} , which contains $\{P\}$ self-conjugately.²⁴

2. *Determination of H_1 .* H_1 is generated by P and some operator Q_1 of G .

$$Q_1^p = P^{hp}.$$

Denoting $Q_1^a P^b Q_1^c P^d \dots$ by the symbol $[a, b, c, d, \dots]$, all operators of H_1 are included in the set $[y, x]; (y = 0, 1, 2, \dots, p-1, x = 0, 1, 2, \dots, p^{m-3} - 1)$.

Since $\{P\}$ is self-conjugate in H_1 ²⁵

$$Q_1^{-1} P Q_1 = P^{1+kp^{m-4}}. \quad (1)$$

²⁴BURNSIDE, *Theory of Groups*, Art. 54, p. 64.

²⁵*Ibid.*, Art. 56, p. 66.

Hence

$$[-y, x, y] = [0, x(1 + kyp^{m-4})] \quad (m > 4). \quad (2)$$

and

$$[y, x]^s = [sy, x \{s + ky \binom{s}{2} p^{m-4}\}]. \quad (3)$$

Placing $y = 1$ and $s = p$ in (3), we have,

$$[Q_1 P^x]^p = Q_1^p P^{xp} = P^{(x+h)p}$$

and if x be so chosen that

$$(x + h) \equiv 0 \pmod{p^{m-4}},$$

$Q = Q_1 P^x$ will be an operator of order p which will be used in place of Q_1 , $Q^p = 1$.

3. *Determination of H_2 .* There is in G a subgroup H_2 of order p^{m-1} , which contains H_1 self-conjugately. H_2 is generated by H_1 , and some operator R_1 of G .

$$R_1^p = P^{lp}.$$

We will now use the symbol $[a, b, c, d, e, f, \dots]$ to denote $R_1^a Q^b P^c R_1^d Q^e P^f \dots$.

The operations of H_2 are given by $[z, y, x]$; ($z, y = 0, 1, \dots, p-1$; $x = 0, 1, \dots, p^{m-3} - 1$). Since H_1 is self-conjugate in H_2

$$R_1^{-1} P R_1 = Q_1^\beta P^{\alpha_1}, \quad (4)$$

$$R_1^{-1} Q R_1 = Q_1^{b_1} P^{\alpha_2 p^{m-4}}. \quad (5)$$

From (4), (5) and (3)

$$[-p, 0, 1, p] = \left[0, \frac{\alpha_1^p - b_1^p}{\alpha_1 - b_1} \beta, \alpha_1^p + \theta p^{m-4} \right] = [0, 0, 1],$$

where

$$\theta = \frac{\alpha_1^p \beta k}{2} \frac{\alpha_1^p - 1}{\alpha_1 - 1} + a\beta \left\{ \frac{\alpha_1^p - 1}{\alpha_1 - b_1} p - \frac{\alpha_1^p - b_1^p}{(\alpha_1 - b_1)^2} \right\}.$$

Hence

$$\frac{\alpha_1^p - b_1^p}{\alpha_1 - b_1} \beta \equiv 0 \pmod{p}, \quad \alpha_1^p + \theta p^{m-4} \equiv 1 \pmod{p^{m-3}}, \quad (6)$$

and $\alpha_1^p \equiv 1 \pmod{p^{m-4}}$, or $\alpha_1 \equiv 1 \pmod{p^{m-5}}$ ($m > 5$), $\alpha_1 = 1 + \alpha_2 p^{m-5}$. Equation (4) is replaced by

$$R_1^{-1} P R_1 = Q^\beta P^{1 + \alpha_2 p^{m-5}}, \quad (7)$$

From (5), (7) and (3).

$$[-p, 1, 0, p] = \left[0, b_1^p, a \frac{b_1^p - 1}{b - 1} p^{m-4} \right].$$

Placing $x = lp$ and $y = 1$ in (2) we have $Q^{-1}P^{lp}Q = P^{lp}$, and

$$b_1^p \equiv 1 \pmod{p}, \quad a \frac{b_1^p - 1}{b_1 - 1} \equiv 0 \pmod{p}.$$

Therefore, $b_1 = 1$.

Substituting 1 for b_1 and $1 + \alpha_2 p^{m-5}$ for α_1 in congruence (6) we find

$$(1 + \alpha_2 p^{m-5})^p \equiv 1 \pmod{p^{m-3}}, \quad \text{or} \quad \alpha_2 \equiv 0 \pmod{p}.$$

Let $\alpha_2 = \alpha p$ and equations (7) and (5) are replaced by

$$R_1^{-1} P R_1 = Q^\beta P^{1+\alpha p^{m-4}}, \quad (8)$$

$$R_1^{-1} Q R_1 = Q P^{\alpha p^{m-4}}. \quad (9)$$

From (8), (9) and (3)

$$[-y, 0, x, y] = \left[0, \beta xy, x + \{ \alpha xy + a\beta x \binom{y}{2} + \beta ky \binom{x}{2} \} p^{m-4} \right], \quad (10)$$

$$[-y, x, 0, y] = [0, x, axy p^{m-4}]. \quad (11)$$

From (2), (10), and (11)

$$[z, y, x]^s = [sz, sy + U_s, sx + V_s p^{m-4}], \quad (12)$$

where

$$\begin{aligned} U_s &= \beta \binom{s}{2} xz, \\ V_s &= \binom{s}{2} \{ \alpha xz + kxy + ayz + \beta k \binom{x}{2} z \} \\ &\quad + \beta k \binom{s}{3} x^2 z + \frac{1}{2} a \beta \binom{s}{2} \left\{ \frac{1}{3!} (2s-1)z - 1 \right\} xz. \end{aligned}$$

Placing $z = 1$, $y = 0$, and $s = p$ in (12)²⁶

$$[R_1 P^x]^p = R_1^p P^{xp} = P^{(x+l)p}.$$

If x be so chosen that

$$x + l \equiv 0 \pmod{p^{m-4}}$$

then $R = R_1 P^x$ is an operator of order p which will be used in place of R_1 , and $R^p = 1$.

²⁶The terms of the form $(Ax + Bx^2)p^{m-4}$ which appear in the exponent of P for $p = 3$ do not alter the conclusion for $m > 5$.

4. *Determination of G.* G is generated by H_2 and some operation S_1 .

$$S_1^p = P^{\lambda p}.$$

Denoting $S_1^a R^b Q^c P^d \dots$ by the symbol $[a, b, c, d, \dots]$ all the operators of G are given by

$$[v, z, y, x]; (v, z, y = 0, 1, \dots, p-1; x = 0, 1, \dots, p^{m-3} - 1).$$

Since H_2 is self-conjugate in G

$$S_1^{-1} P S_1 = R^\gamma Q^s P^{\epsilon_1}, \quad (13)$$

$$S_1^{-1} Q S_1 = R^c Q^d P^{\epsilon p^{m-4}}, \quad (14)$$

$$S_1 R S_1 = R^f Q^g P^{j p^{m-4}}. \quad (15)$$

From (13), (14), (15), and (12)

$$[-p, 0, 0, 1, p] = [0, L, M, \epsilon_1^p + N p^{m-4}] = [0, 0, 0, 1]$$

and

$$\epsilon_1^p \equiv 1 \pmod{p^{m-4}} \quad \text{or} \quad \epsilon_1 \equiv 1 \pmod{p^{m-5}} \quad (m > 5).$$

Let $\epsilon_1 = 1 + \epsilon_2 p^{m-5}$. Equation (13) is now replaced by

$$S_1^{-1} P S_1 = R^\gamma Q^\delta P^{1+\epsilon_2 p^{m-5}}. \quad (16)$$

If $\lambda = 0 \pmod{p}$ and $\lambda = \lambda' p$,

$$\begin{aligned} [1, 0, 0, 1]^p &= [p, 0, 0, p + \epsilon \binom{p}{2} p^{m-5} + W p^{m-4}] \\ &= [0, 0, 0, p + \lambda' p^2 + W' p^{m-4}] \end{aligned}$$

and for $m > 5$ $S_1 P$ is of order p^{m-3} . We will take this in place of S_1 and assume $dv[\lambda, p] = 1$.

$$S_1^{p^{m-3}} = 1.$$

There is in G a subgroup H'_1 of order p^{m-2} which contains $\{S_1\}$ self-conjugately. $H'_1 = \{S_1, S_1^y R^z Q^y P^x\}$ and the operator $T = R^z Q^y P^x$ is in H'_1 .

There are two cases for discussion.

1°. Where x is prime to p .

T is an operator of H_2 of order p^{m-3} and will be taken as P . Then

$$H'_1 = \{S_1, P\}.$$

Equation (16) becomes

$$S_1^{-1} P S_1 = P^{1+\epsilon p^{m-4}}.$$

There is in G a subgroup H'_2 of order p^{m-1} which contains H'_1 self-conjugately.

$$H'_2 = \{H'_1, S_1^{v'} R^{z'} Q^{y'} P^{x'}\}.$$

$T' = R^{z'} Q^{y'}$ is in H'_2 and also in H_2 and is taken as Q , since $\{P, T'\}$ is of order p^{m-2} .

$H'_2 = \{H'_1, Q\} = \{S_1, H_1\}$ and in this case c may be taken $\equiv 0 \pmod{p}$.

2°. Where $x = x_1 p$. P^p is in $\{S_1\}$ since λ is prime to p . In the present case $R^z Q^y$ is in H'_1 and also in H_2 . If $z \not\equiv 0 \pmod{p}$ take $R^z Q^y$ as R ; if $z \equiv 0 \pmod{p}$ take it as Q .

$$H'_1 = \{S_1, R\} \quad \text{or} \quad \{S_1, Q\},$$

and

$$R^{-1} S_1 R = S_1^{1+k'p^{m-4}} \quad \text{or} \quad Q^{-1} S_1 Q = S_1^{1+k''p^{m-4}}.$$

On rearranging these take the forms

$$S_1^{-1} R S_1 = R S_1^{mp^{m-4}} = R P^{jp^{m-4}} \quad \text{or} \quad S_1^{-1} Q S_1 = Q S_1^{n'p^{m-4}} = Q P^{ep^{m-4}},$$

and either c or g may be taken $\equiv 0 \pmod{p}$,

$$cg \equiv 0 \pmod{p}. \quad (17)$$

From (14), (15), (16), (12) and (17)

$$[-p, 0, 1, 0, p] = \left[0, c \frac{d^p - f^p}{d - f}, d^p, W p^{m-4} \right].$$

Place $x = \lambda p$ and $y = 1$ in (12)

$$Q^{-1} P^{\lambda p} Q = P^{\lambda p} \quad \text{or} \quad S_1^p Q S_1^p = Q,$$

and

$$d^p \equiv 1 \pmod{p}, \quad d = 1.$$

Equation (14) is replaced by

$$S_1^{-1} Q S_1 = R^c Q P^{ep^{m-4}}. \quad (18)$$

From (15), (18), (17), (16) and (12)

$$[-p, 1, 0, 0, p] = \left[0, f^p, \frac{d^p - f^p}{d - f} g, W' p^{m-4} \right].$$

Placing $x = \lambda p$, $y = 1$ in (10)

$$R^{-1} P^{\lambda p} R = P^{\lambda p},$$

and $f^p \equiv 1 \pmod{p}$, $f = 1$. Equation (15) is replaced by

$$S_1^{-1} R S_1 = R Q^g P^j p^{m-4}. \quad (19)$$

From (16), (18), (19) and (12)

$$S_1^{-p} P S_1^p = P^{1+\epsilon_2 p^{m-4}} = P$$

and $\epsilon_2 \equiv 0 \pmod{p}$. Let $\epsilon_2 = \epsilon p$ and (16) is replaced by

$$S_1^{-1} P S_1 = R^\gamma Q^\delta P^{1+\epsilon p^{m-4}}. \quad (20)$$

Transforming both sides of (1), (8) and (9) by S_1

$$\begin{aligned} S_1^{-1} Q^{-1} S_1 \cdot S_1^{-1} P S_1 \cdot S_1^{-1} Q S_1 &= S_1^{-1} P^{1+k p^{m-4}} S_1, \\ S_1^{-1} R^{-1} S_1 \cdot S_1^{-1} P S_1 \cdot S_1^{-1} R S_1 &= S_1^{-1} Q^\beta S_1 \cdot S_1^{-1} P^{1+\alpha p^{m-4}} S_1, \\ S_1^{-1} R^{-1} S_1 \cdot S_1^{-1} Q S_1 \cdot S_1^{-1} R S_1 &= S_1^{-1} Q S_1 \cdot S_1^{-1} P^{a p^{m-4}} S_1. \end{aligned}$$

Reducing these by (18), (19), (20) and (12) and rearranging

$$\begin{aligned} &[0, \gamma, \delta + \beta c, 1 + \{\epsilon + \alpha c + k + a c \delta + a \beta \binom{c}{2} - a \gamma\} p^{m-4}] \\ &= [0, \gamma, \delta, 1 + (\epsilon + k) p^{m-4}]. \\ &[0, \gamma, \beta + \delta, 1 + \{k g + \epsilon + \alpha + a \delta - a \gamma g\} p^{m-4}] \\ &= [0, \gamma + \beta c, \beta + \delta, 1 + \{\epsilon + \alpha + \beta e + \alpha \binom{\beta}{2} c + a \beta \gamma\} p^{m-4}], \\ &[0, c, 1, (e + a) p^{m-4}] = [0, c, 1, (e + a) p^{m-4}]. \end{aligned}$$

The first gives

$$\beta c \equiv 0 \pmod{p}, \quad (21)$$

$$a c + a c \delta - a \gamma \equiv 0 \pmod{p}. \quad (22)$$

Multiplying this last by g

$$a g \gamma \equiv 0 \pmod{p}. \quad (23)$$

From the second equation above

$$g k + \alpha \delta \equiv \beta e + a \beta \gamma \pmod{p}. \quad (24)$$

Multiplying by c

$$a c \delta \equiv 0 \pmod{p}. \quad (25)$$

These relations among the constants *must be satisfied* in order that our equations should define a group.

From (20), (19), (18) and (12)

$$[-y, 0, 0, x, y] = [0, \gamma xy + \chi_1(x, y), \delta xy + \phi_1(x, y), x + \Theta_1(x, y)p^{m-4}], \quad (26)$$

$$[-y, 0, x, 0, y] = [0, cxy, x, \Theta_2(x, y)p^{m-4}], \quad (27)$$

$$[-y, x, 0, 0, y] = [0, x, gxy, \Theta_3(x, y)p^{m-4}], \quad (28)$$

where

$$\chi_1(x, y) = c\delta x \binom{y}{2},$$

$$\phi_1(x, y) = \gamma gx \binom{y}{2} + \beta \gamma \binom{x}{2} y,$$

$$\begin{aligned} \Theta_1(x, y) = \epsilon xy + \binom{y}{2} [\gamma jx + e\delta x + a\delta\gamma + (\alpha\gamma + k\delta) \binom{x}{2}] \\ + \binom{y}{3} [c\delta j + eg\gamma]x + \binom{x}{2} [\alpha\gamma y + \delta ky + a\delta\gamma y^2] + \beta\gamma k \binom{x}{3} y^2, \end{aligned}$$

$$\Theta_2(x, y) = \epsilon xy + cjx \binom{y}{2} + ac \binom{x}{2} y,$$

$$\Theta_3(x, y) = jxy + egx \binom{y}{2} + ag \binom{x}{2} y.$$

Let a general power of any operator be

$$[v, z, y, x]^s = [sv, sz + U_s, sy + V_s, sx + W_s p^{m-4}]. \quad (29)$$

Multiplying both sides by $[v, z, y, x]$ and reducing by (2), (10), (11), (26), (27) and (28), we find

$$U_{s+1} \equiv U_s + (cy + \gamma x)sv + c\delta \binom{sv}{2} x \pmod{p},$$

$$V_{s+1} \equiv V_s + (gz + \delta x)sv + \gamma g \binom{sv}{2} x + \beta \gamma \binom{x}{2} sv + \beta (sz + U_s)x \pmod{p},$$

$$\begin{aligned} W_{s+1} \equiv W_s + \Theta_1(x, sv) + \{ey + jz + a\gamma xy + ac \binom{y}{2} + ag \binom{z}{2}\} sv \\ + \{\alpha x + \beta k \binom{x}{2} + ay + a\delta sx + \alpha gsvz\} sz + ksxy \\ + \binom{sv}{2} \{c jy + egz\} + U_s \{\alpha x + \beta k \binom{x}{2} + ay + a(\delta x + gz)sv\} \\ + a\beta \binom{sz+U_s}{2} x + kV_s x \pmod{p}. \end{aligned}$$

From (29)

$$U_1 \equiv 0, \quad V_1 \equiv 0, \quad W_1 \equiv 0 \pmod{p}.$$

A continued use of the above congruences give

$$U_s \equiv (cy + \gamma x) \binom{s}{2} v + \frac{1}{2} c\delta xv \left\{ \frac{1}{3} (2s-1)v - 1 \right\} \binom{s}{2} \pmod{p},$$

$$\begin{aligned} V_s \equiv \{gz + \delta x + \beta \gamma \binom{x}{2} v + \beta xz\} \binom{s}{2} \\ + \frac{1}{2} \gamma gxv \left\{ \frac{1}{3} (2s-1)v - 1 \right\} \binom{s}{2} + \beta \gamma \binom{s}{3} x^2 v \pmod{p}, \end{aligned}$$

$$\begin{aligned}
W_s \equiv \binom{s}{2} & \left\{ \epsilon xv + egv + (\alpha\gamma + \delta kv + \beta kz) \binom{s}{2} + \beta\gamma k \binom{x}{3} v + ac \binom{y}{2} v \right. \\
& + jvz + ag \binom{z}{2} v + \alpha xz + kxy + a\gamma xyv + ayz \left. \right\} + \binom{s}{3} \left\{ \alpha cxyv \right. \\
& + \alpha\gamma x^2 v + 2\beta\gamma k \binom{x}{2} xv + gkxzv + \delta kx^2 v + \beta kx^2 z + acvy^2 \\
& + a\gamma xvy \left. \right\} + \beta k\gamma \binom{s}{4} x^3 v + \binom{s}{2} \frac{2s-1}{3} \left\{ a\delta\gamma \binom{x}{2} v^2 + a\delta xzv \right. \\
& + agvz^2 \left. \right\} + \frac{1}{2} v \binom{s}{2} \left\{ \frac{1}{3} (2s-1)v - 1 \right\} \left\{ \gamma jx + e\delta x + a\delta\gamma x \right. \\
& + \alpha c\delta \binom{x}{2} + \gamma gk \binom{x}{2} + c jy + egz \left. \right\} + \frac{1}{6} \binom{s}{2} \left\{ \binom{s}{2} v^2 - (2s-1)v \right. \\
& + 2 \left. \right\} \left\{ c\delta jx + eg\gamma x \right\} v + \frac{1}{2} \binom{s}{3} \left\{ \frac{1}{2} (s-1)v - 1 \right\} \left\{ \alpha c\delta \right. \\
& + \gamma gk \left. \right\} x^2 v + \frac{1}{2} a\beta x \binom{s}{2} \left\{ \frac{1}{3} (2s-1)z - 1 \right\} z \\
& + \frac{1}{2} a\delta\gamma x^2 v \binom{s}{3} \frac{1}{2} (3s-1) \pmod{p}
\end{aligned}$$

Placing $v = 1$, $z = y = s = p$ in (29)²⁷

$$[S_1 P^x]^p = S_1^p P^{xp} = P^{(\lambda+x)p} \quad (p > 3).$$

If x be so chosen that

$$x + \lambda \equiv 0 \pmod{p^{m-4}}.$$

$S = S_1 P^x$ is an operator of order p and is taken in place of S_1 .

$$S^p = 1.$$

The substitution of S for S_1 leaves congruence (17) invariant.

5. *Transformation of the groups.* All groups of this class are given by

$$G : \begin{cases} Q^{-1} P Q = P^{1+kp^{m-4}}, \\ R^{-1} P R = Q^\beta P^{1+\alpha p^{m-4}}, \\ R^{-1} Q R = Q P^{\alpha p^{m-4}}, \\ S^{-1} P S = R^\gamma Q^\delta P^{1+\epsilon p^{m-4}}, \\ S^{-1} Q S = R^c Q P^{\epsilon p^{m-4}}, \\ S^{-1} R S = R Q^g P^{j p^{m-4}}, \end{cases} \quad (30)$$

with

$$P^{p^{m-3}} = 1, \quad Q^p = R^p = S^p = 1,$$

²⁷For $p = 3$ and $c\delta \equiv \gamma g \equiv \beta\gamma \equiv 0 \pmod{p}$ there are terms of the form $(A + Bx + Cx^2 + Dx^3)p^{m-4}$ in the exponent of P . For $m > 5$ these do not vitiate our conclusion. For $p = 3$ and $c\delta, \gamma g$, or $\beta\gamma$ prime to p , $[S_1 P^x]^p$ is not contained in $\{P\}$ and the groups defined belong to Class II.

($k, \beta, \alpha, a, \gamma, \delta, \epsilon, c, e, g, j = 0, 1, 2, \dots, p-1$).

These constants are however subject to conditions (17), (21), (22), (23), (24) and (25). Not all these groups are distinct. Suppose that G and G' of the above set are simply isomorphic and that the correspondence is given by

$$C = \begin{bmatrix} S, & R, & Q, & P \\ S'_1, & R'_1, & Q'_1, & P'_1 \end{bmatrix}.$$

Inspection of (29) gives

$$\begin{aligned} S'_1 &= S'^{v''''} R'^{z''''} Q'^{y''''} P'^{x''''} p^{m-4}, \\ R'_1 &= S'^{v'''} R'^{z'''} Q'^{y'''} P'^{x'''} p^{m-4}, \\ Q'_1 &= S'^{v''} R'^{z''} Q'^{y''} P'^{x''} p^{m-4}, \\ P'_1 &= S'^v R'^z Q'^y P'^x, \end{aligned}$$

in which x and one out of each of the sets $v', z', y', x'; v'', z'', y'', x''; v''', z''', y''', x'''; v'''' , z'''' , y'''' , x''''$ are prime to p .

Since $S, R, Q,$ and P satisfy equations (30), S'_1, R'_1, Q'_1 and P'_1 also satisfy them. Substituting these operators and reducing in terms of $S', R', Q',$ and P' we get the six equations

$$[V'_\kappa, Z'_\kappa, Y'_\kappa, X'_\kappa] = [V_\kappa, Z_\kappa, Y_\kappa, X_\kappa] \quad (\kappa = 1, 2, 3, 4, 5, 6), \quad (31)$$

which give the following twenty-four congruences

$$\begin{cases} V'_\kappa \equiv V_\kappa \pmod{p}, \\ Z'_\kappa \equiv Z_\kappa \pmod{p}, \\ Y'_\kappa \equiv Y_\kappa \pmod{p}, \\ X'_\kappa \equiv X_\kappa \pmod{p^{m-3}}, \end{cases} \quad (32)$$

where

$$\begin{aligned} V'_1 &= v, \quad V_1 = v, \\ Z'_1 &= Z + c'(yv' - y'v) + \gamma'xv' + c\delta x \binom{v'}{2}, \quad Z_1 = z, \\ Y'_1 &= y + g'(zv' - z'v) + \delta'xv' + \gamma'g'x \binom{v'}{2} + \beta'xz', \quad Y_1 = y, \\ X'_1 &= x + \left\{ \epsilon'xv' + (\gamma'j'x + e'\delta'x + a'\delta'\gamma'x) \binom{v'}{2} + c'\delta'j' \binom{v'}{3} + (\alpha'\gamma'v' + \delta'k'v' \right. \\ &\quad + a'\delta'\gamma'v^2 + \beta'k'z') \binom{x}{2} + j'(zv' - z'v) + e'g'[z \binom{v'}{2} - z' \binom{v'}{2}] \\ &\quad + a'g'[(\binom{z}{2}v' + \binom{z'}{2}v - zz'v) + e'(yv' - y'v) + c'j'[y \binom{v'}{2} - y' \binom{v'}{2}]] \\ &\quad + a'c'[(\binom{y}{2}v' + v \binom{-y'}{2}) - yy'v] + a'(yz' - y'z) - a'\beta'xz'^2 + \alpha'xz' \\ &\quad \left. + \alpha'\beta'x \binom{z'}{2} + a'\gamma'x(y - y')v' + k'xy' \right\} p^{m-4}, \\ X_1 &= x + kxp^{m-4}, \end{aligned}$$

$$\begin{aligned}
V_2' &= v, \quad V_2 = v + \beta v', \\
Z_2' &= z + c'(yv'' - y''v) + \gamma' xv'' + e'\delta'(v_2''), \quad Z_2 = z + \beta z' + c'\beta y'v, \\
Y_2' &= y + g'(zv'' - z''v) + \delta' xv'' + \gamma' g'x(v_2'') + \beta' \gamma' \binom{x}{2} v'' + \beta' xz'', \\
Y_2 &= y + \beta y' + g'\beta z'v, \\
X_2' &= x + \left\{ \Theta_1'(x, v'') + j'(zv'' - z''v) + e'g'[z(v_2'') - z''(v_2')] + a'g'[\binom{x}{2}v'' \right. \\
&\quad + \binom{-z''}{2}v - zz''v] + e'(yv'' - y''v) + c'j'[y(v_2'') - y''(v_2')] + a'c'[\binom{y}{2}v'' \\
&\quad + \binom{-y''}{2}v - yy''v''] + a'g'(zv'' - z''v)z'' + a'(yz'' - y''z) + a'\delta'v''z'' \\
&\quad \left. + a'\gamma'(y - y'')v''x + \alpha'xz'' + a'\beta'x(z_2'') + \beta'k'\binom{x}{2}z'' + k'xy'' \right\} p^{m-4}, \\
X_2 &= x + \left\{ \alpha x + \beta x' + a'\binom{\beta}{2}y'z' + e'\beta v y' + (c'j'\beta + e'g'\beta z')\binom{v}{2} \right. \\
&\quad \left. + a'c'\binom{\beta y'}{2}v + j'\beta v z' + a'g'\binom{\beta z'}{2} + a'\beta(g'z'v + y'z) \right\} p^{m-4}, \\
V_3' &= v', \quad V_3 = v', \\
Z_3' &= z' + c'(y'v'' - y''v'), \quad Z_3 = z', \\
Y_3' &= y' + g'(z'v'' - z''v'), \quad Y_3 = y', \\
X_3' &= \left\{ x' + j'(z'v'' - z''v') + e'g'[(v_2'')z' - (v_2')z''] + a'g'[\binom{z'}{2}v'' + \binom{-z''}{2}v' \right. \\
&\quad - z'z''v'] + e'(y'v'' - y''v') + c'j'[y'(v_2'') - y''(v_2')] + a'c'[\binom{y'}{2}v'' + \binom{-y''}{2}v' \\
&\quad \left. - y''y'v''] + a'(y'z'' - y''z') \right\} p^{m-4}, \\
X_4 &= (x' + a'x)p^{m-4}, \\
V_4' &= v, \quad V_4 = v + \gamma v'' + \delta v', \\
Z_4' &= z + c'(yv''' - y'''v) + \gamma' xv''' + c'\delta'x(v_2'''), \\
Z_4 &= z + \gamma z'' + \delta z' + c'[\binom{\gamma}{2}v''y'' + \binom{\delta}{2}v'y'] + c'(\gamma y'' + \delta y')v + c'\gamma \delta y''v, \\
Y_4' &= y + g'(zv''' - z'''v) + \delta' xv''' + \gamma' g'x(v_2''') + \beta' \gamma' \binom{x}{2} v''' + \beta' xz''', \\
Y_4 &= y + \gamma y'' + \delta y' + g'[\binom{\gamma}{2}v''z'' + \binom{\delta}{2}v'z'] + g'(\gamma z'' + \delta z')v + g'\delta \gamma v'z'', \\
X_4' &= x + \left\{ \Theta_1'(x, v''') + j'(zv''' - z'''v) + e'g'[(v_2''')z - \binom{v}{2}z'''] + a'g'[\binom{z}{2}v''' \right. \\
&\quad + \binom{-z'''}{2}v - zz'''v] + e'(yv''' - y'''v) + c'j'[y(v_2''') - y'''(v_2')] \\
&\quad + a'c'[\binom{y}{2}v''' + \binom{-y'''}{2}v - yy'''v'''] + a'g'(v'''z - vz''')z''' \\
&\quad + a'(yz''' - y'''z) + a'\delta'xz'''v''' + a'\gamma'x(y - y''')v''' + \alpha'xz''' \\
&\quad \left. + a'\beta'x(z_2''') + \beta'k'z'''(x) + k'xy''' \right\} p^{m-4}.
\end{aligned}$$

$$\begin{aligned}
X_4 = & x + \left\{ \epsilon x + \delta x' + \gamma x'' + \binom{\gamma}{2} [a'c' \binom{y''}{2} v'' + a'y''z'' + e'v''y'' + j'v''z'' \right. \\
& + a'g' \binom{z''}{2} v'' + (c'j'v''y'' + e'g'v''z'')(v + \delta v') + a'(z + \delta z')v''z'' \\
& + \frac{2\gamma - 1}{3} a'g'v''z''^2 + \frac{1}{2} \left[\frac{1}{3} (2\gamma - 1)v'' - 1 \right] (c'j'y'' + e'g'z'')v'' \\
& + \binom{\gamma}{3} a'c'v''y'' + \binom{\delta}{2} [a'c' \binom{y'}{2} v' + a'y'z' + e'v'y' + j'v'z' \\
& + a'g' \binom{z'}{2} v' + j'c'vv'y' + e'g'vv'z' + a'g'v'zz' + a'c'\gamma y'y'v' \\
& + \frac{2\delta - 1}{3} a'g'v'z'^2 + \frac{1}{2} \left\{ \frac{1}{3} (2\delta - 1)v' - 1 \right\} (c'j'y' + e'g'z') \\
& + \binom{\delta}{3} a'c'v'y'^2 + (v + \delta v') [j'\gamma z'' + \binom{\gamma z''}{2} a'g' + e'\gamma y'' + \binom{\gamma y''}{2} a'c' \\
& + a'g'(z + \delta z')] + \binom{v + \delta v'}{2} [e'g'\gamma z'' + c'j'\gamma y''] + \delta [(e'g'z' \\
& + c'j'y') \binom{v}{2} + e'vy' + j'z' + a'zy + a'g'vz'z' + a'\gamma z'y'' + a'c'\gamma vy'y''] \\
& \left. + a'g' \binom{\delta z'}{2} v + a'c' \binom{\delta y'}{2} v + a'\gamma zy'' \right\} p^{m-4},
\end{aligned}$$

$$V'_5 = v', \quad V_5 = v' + cv'',$$

$$Z'_5 = z' + c'(y'v''' - y'''v'), \quad Z_5 = z' + cz'' + c'cy''v,$$

$$Y'_5 = y' + g'(z'v''' - z'''v'), \quad Y_5 = y' + cy'' + g'cv'z'',$$

$$\begin{aligned}
X'_5 = & \left\{ x' + j'(z'v''' - z'''v') + e'g' \left[\binom{v'''}{2} z' - \binom{v'}{2} z'''' \right] + a'g' \left[\binom{z'}{2} v'''' \right. \right. \\
& + \binom{-z'''}{2} v' - z'z''''v'] + c'(y'v''' - y'''v') + c'j' [y' \binom{\delta'''}{2} - y''' \binom{v'}{2}] \\
& \left. + a'c' \left[\binom{y''}{2} v'''' + \binom{-y'''}{2} v' - y'y''''v'''' \right] + a'(y'z'''' - y''''z') \right\} p^{m-4},
\end{aligned}$$

$$\begin{aligned}
X_5 = & \left\{ x' + \epsilon x + cx'' + a' \binom{c}{2} y''z'' + j'cv'z'' + (e'g'cz'' + c'cj'y'') \binom{v'}{2} + e'cy''v \right. \\
& \left. + a'cy''z' + a'g'z'v' + a'g' \binom{cz''}{2} + a'c' \binom{cy''}{2} \right\} p^{m-4},
\end{aligned}$$

$$V'_6 = v'', \quad V_6 = v'' + gv',$$

$$Z'_6 = z'' + c'(y''v'''' - y''''v''), \quad Z_6 = z'' + gz',$$

$$Y'_6 = y'' + g'(z''v'''' - z''''v''), \quad Y_6 = y'' + gy',$$

$$\begin{aligned}
X'_6 = & \left\{ x'' + j'(z''v'''' - z''''v'') + e'g' \left[\binom{v''''}{2} z'' - \binom{v''}{2} z'''' \right] + a'g' \left[\binom{z''}{2} v'''' \right. \right. \\
& + \binom{-z''''}{2} v'' - z''z''''v'''] + e'(y''v'''' - y''''v'') + c'j' [y'' \binom{v''''}{2} - y'''' \binom{v''}{2}] \\
& \left. + a'c' \left[\binom{y''}{2} v'''' + \binom{-y''''}{2} v'' - y''y''''v'''' \right] + a'(y''z'''' - y''''z'') \right\} p^{m-4},
\end{aligned}$$

$$X_6 = \{x'' + jx + gx' + a'gy''z'\} p^{m-4}.$$

The necessary and sufficient condition for the simple isomorphism of the two groups G and G' is that congruences (32) shall be consistent and admit of solution subject to conditions derived below.

6. *Conditions of transformation.* Since Q is not contained in $\{P\}$, R is not contained in $\{Q, P\}$, and S is not contained in $\{R, Q, P\}$, then Q'_1 is not

contained in $\{P'_1\}$, R'_1 is not contained in $\{Q'_1, P'_1\}$, and S'_1 is not contained in $\{R'_1, Q'_1, P'_1\}$.

Let

$$Q_1^{s'} = P_1^{s'p^{m-4}}.$$

This equation becomes in terms of S' , R' , Q' and P'

$$[s'v', s'z' + c' \binom{s'}{2} v'y', s'y' + g' \binom{s'}{2} v'z', Dp^{m-4}] = [0, 0, 0, sxp^{m-4}],$$

and

$$s'v' \equiv s'z' \equiv s'y' \equiv 0 \pmod{p}.$$

At least one of the three quantities v' , z' or y' is prime to p , since otherwise s' may be taken = 1.

Let

$$R_1^{s''} = Q_1^{s'} P_1^{s'p^{m-4}},$$

or in terms of S' , R' , Q' and P'

$$\begin{aligned} [s''v'', s''z'' + c' \binom{s''}{2} v''y'', s''y'' + g' \binom{s''}{2} v''z'', Ep^{m-4}] \\ = [s'v', s'z' + c' \binom{s'}{2} v'y', s'y' + g' \binom{s'}{2} v'z', E_1p^{m-4}], \end{aligned}$$

and

$$\begin{aligned} s''v'' &\equiv s'v' \pmod{p}, \\ s''z'' + c' \binom{s''}{2} v''y'' &\equiv s'z' + c' \binom{s'}{2} v'y' \pmod{p}, \\ s''y'' + g' \binom{s''}{2} v''z'' &\equiv s'y' + g' \binom{s'}{2} v'z' \pmod{p}. \end{aligned}$$

Since $c'g' \equiv 0 \pmod{p}$, suppose $g' \equiv 0 \pmod{p}$. Elimination of s' between the last two give by means of the congruence $Z'_3 \equiv Z_3 \pmod{p}$,

$$s''\{2(y'z'' - y''z') + c'y'y''(v' - v'')\} \equiv 0 \pmod{p},$$

between the first two

$$s''\{2(v'z'' - v''z') + c'v'v''(y' - y'')\} \equiv 0 \pmod{p},$$

and between the first and last

$$s''(y'v'' - y''v') \equiv 0 \pmod{p}.$$

At least one of the three above coefficients of s'' is prime to p , since otherwise s'' may be taken = 1.

Let

$$S_1^{s'''} = R_1^{s''} Q_1^{s'} P_1^{s'p^{m-4}}$$

or, in terms of S' , R' , Q' , and P'

$$\begin{aligned} & [s'''v''', s'''z''' + c' \binom{s'''}{2} v'''y''', s'''y''' + g' \binom{s'''}{2} v'''z''', E_2p^{m-4}] \\ & = [s''v'' + s'v', s''z'' + s'z' + c' \{ \binom{s''}{2} v''y'' + \binom{s'}{2} v'y' + s's''y''v' \}, \\ & \quad s''y'' + s'y' + g' \{ \binom{s''}{2} v''z'' + \binom{s'}{2} v'z' + s's''v'z'' \}, E_3p^{m-4}] \end{aligned}$$

and

$$\begin{aligned} s'''v''' & \equiv s''v'' + s'v' \pmod{p}, \\ s'''z''' + c' \binom{s'''}{2} v'''y''' & \equiv s''z'' + s'z' + c' \{ \binom{s''}{2} v''y'' + \binom{s'}{2} v'y' + s's''y''v' \} \pmod{p}, \\ s'''y''' + g' \binom{s'''}{2} v'''z''' & \equiv s''y'' + s'y' + g' \{ \binom{s''}{2} v''z'' + \binom{s'}{2} v'z' + s's''z''v' \} \pmod{p}. \end{aligned}$$

If $g' \equiv 0$ and $c' \not\equiv 0 \pmod{p}$ the congruence $Z'_3 \equiv Z_3 \pmod{p}$ gives

$$(y'v'' - y''v') \equiv 0 \pmod{p}.$$

Elimination in this case of s'' between the first and last congruences gives

$$s'''(y''v''' - y'''v'') \equiv 0 \pmod{p}.$$

Elimination of s'' between the first and second, and between the second and third, followed by elimination of s' between the two results, gives

$$s''' \left(z''^2 - c'y''z''v' + \frac{c'^2}{4} y''v'' \right) (y'v''' - y'''v') \equiv 0 \pmod{p}.$$

Either $(y''v''' - y'''v'')$, or $(y'v''' - y'''v')$ is prime to p , since otherwise s''' may be taken = 1.

A similar set of conditions holds for $c' \equiv 0$ and $g' \not\equiv 0 \pmod{p}$.

When $c' \equiv g' \equiv 0 \pmod{p}$ elimination of s' and s'' between the three congruences gives

$$s''' \Delta \equiv s''' \begin{vmatrix} v' & v'' & v''' \\ y' & y'' & y''' \\ z' & z'' & z''' \end{vmatrix} \equiv 0 \pmod{p}$$

and Δ is prime to p , since otherwise s''' may be taken = 1.

7. *Reduction to types.* In the discussion of congruences (32), the group G' is taken from the simplest case and we associate with it all simply isomorphic groups G .

I.

A.

B.

	a_2	β_2	c_2	g_2	γ_2	δ_2		k_2	α_2	ϵ_2	e_2	j_2
1	1	1	1	1	1	1	1	1	1	1	1	1
2	0	1	1	1	1	1	2	0	1	1	1	1
3	0	0	1	1	1	1	3	1	0	1	1	1
4	0	0	1	1	1	0	4	1	1	0	1	1
5	0	0	1	0	1	1	5	1	1	1	0	1
6	0	0	1	0	1	0	6	1	1	1	1	0
7	0	1	0	1	1	1	7	0	0	1	1	1
8	0	1	0	1	0	1	8	0	1	0	1	1
9	0	1	1	0	1	1	9	0	1	1	0	1
10	0	1	1	0	1	0	10	0	1	1	1	0
11	1	0	1	1	1	1	11	1	0	0	1	1
12	1	0	1	0	1	1	12	1	0	1	0	1
13	1	0	1	1	0	1	13	1	0	1	1	0
14	1	0	1	1	1	0	14	1	1	0	0	1
15	1	0	1	0	0	1	15	1	1	0	1	0
16	1	0	1	0	1	0	16	1	1	1	0	0
17	1	0	1	1	0	0	17	0	0	0	1	1
18	1	0	1	0	0	0	18	0	0	1	0	1
19	1	1	0	1	1	1	19	0	0	1	1	0
20	1	1	0	1	0	1	20	0	1	0	0	1
21	1	1	0	1	1	0	21	0	1	0	1	0
22	1	1	0	1	0	0	22	0	1	1	0	0
23	1	1	1	0	1	1	23	1	0	0	0	1
24	1	1	1	1	0	1	24	1	0	0	1	0
25	1	1	1	1	1	0	25	1	0	1	0	0
26	1	1	1	0	0	1	26	1	1	0	0	0
27	1	1	1	0	1	0	27	0	0	0	0	1
28	1	1	1	1	0	0	28	0	0	0	1	0
29	1	1	1	0	0	0	29	0	0	1	0	0
							30	0	1	0	0	0
							31	1	0	0	0	0
							32	0	0	0	0	0

II.

A.

	1	2	3	4	5	6	7	8	9	10
1	×	×	×		19 ₆		19 ₆		19 ₆	
2	×	2 ₁	3 ₁			19 ₆	19 ₆			19 ₆
3	1 ₂	2 ₁	3 ₁		19 ₆			19 ₆	19 ₆	
4	1 ₂	×	×		19 ₆		19 ₆		19 ₆	
5	2 ₁	2 ₁		*		19 ₆	19 ₆		19 ₆	
6	2 ₁	2 ₁	3 ₁		19 ₆		19 ₆		19 ₆	
7	1 ₂	2 ₁	3 ₁			19 ₆		19 ₆		19 ₆
8	1 ₂	2 ₄	3 ₄			19 ₆	19 ₆			19 ₆
9	2 ₁	2 ₁		*	19 ₆	19 ₆	19 ₆			19 ₆
10	2 ₄	2 ₄	3 ₄			19 ₆	19 ₆			19 ₆
11	1 ₂	2 ₄	3 ₄		19 ₆			19 ₆	19 ₆	
12	2 ₄	2 ₄		*		19 ₆		19 ₆	19 ₆	
13	2 ₁	2 ₁	3 ₁		19 ₆			19 ₆	19 ₆	
14	2 ₁	2 ₄		*		19 ₆	19 ₆		19 ₆	
15	2 ₁	2 ₄	3 ₄		19 ₆		19 ₆		19 ₆	
16	2 ₁	2 ₁		*		19 ₆	19 ₆		19 ₆	
17	1 ₂	2 ₄	3 ₄			19 ₆		19 ₆		19 ₆
18	2 ₄	2 ₄		*	19 ₆	19 ₆		19 ₆		19 ₆
19	2 ₄	2 ₄	3 ₄			19 ₆		19 ₆		19 ₆
20	2 ₁	2 ₄		*	19 ₆	19 ₆	19 ₆			19 ₆
21	2 ₄	*	*			19 ₆	19 ₆			19 ₆
22	2 ₄	2 ₄		*	19 ₆	19 ₆	19 ₆			19 ₆
23	2 ₄	*		*		19 ₆		19 ₆	19 ₆	
24	2 ₁	2 ₄	3 ₄			19 ₆			19 ₆	19 ₆
25	2 ₄	2 ₄		*		19 ₆		19 ₆	19 ₆	
26	2 ₁	2 ₄		*		19 ₆	19 ₆		19 ₆	
27	2 ₄	*		*	19 ₆	19 ₆		19 ₆		19 ₆
28	2 ₄	*	*			19 ₆		19 ₆		19 ₆
29	*	*		*	19 ₆	19 ₆		19 ₆		19 ₆
30	2 ₄	*		*	19 ₆	19 ₆	19 ₆			19 ₆
31	2 ₄	*		*		19 ₆		19 ₆	19 ₆	
32	*	*		*	19 ₆	19 ₆		19 ₆		19 ₆

B.

II. (continued)

A.

	11	12	13	14	15	16	17	18	19
1	×	19 ₁	×	11 ₁	19 ₁	19 ₁	13 ₁	19 ₁	×
2	25 ₂		×	25 ₂			13 ₂		×
3	11 ₁	19 ₂	13 ₁	24 ₂	21 ₂	19 ₂	13 ₁	21 ₂	
4	24 ₂	19 ₂	13 ₁	24 ₂	19 ₁	19 ₂	13 ₁	19 ₁	19 ₂
5									19 ₁
6	×	19 ₂	×	11 ₆	21 ₂	19 ₂	13 ₆	21 ₂	×
7	25 ₂		13 ₂	25 ₂			13 ₂		
8	25 ₂		13 ₂	25 ₂			13 ₂		19 ₂
9		19 ₆			21 ₆	19 ₆		21 ₆	19 ₂
10	25 ₁₀		×	25 ₁₀			13 ₁₀		19 ₆
11	24 ₂	19 ₂	13 ₁	*	21 ₂	19 ₂	13 ₁	21 ₂	
12									
13	11 ₆	*	13 ₆	11 ₆	*	19 ₂	13 ₆	*	
14									19 ₂
15	11 ₆	19 ₂	13 ₆	11 ₆	21 ₂	19 ₂	13 ₆	21 ₂	19 ₆
16									19 ₆
17	25 ₂		13 ₂	25 ₂			13 ₂		
18		19 ₆			21 ₆	19 ₆		21 ₆	
19	25 ₁₀		13 ₁₀	25 ₁₀			13 ₁₀		
20		19 ₆			21 ₆	19 ₆		21 ₆	19 ₂
21	25 ₁₀		13 ₁₀	25 ₁₀			13 ₁₀		19 ₆
22		19 ₆			21 ₆	19 ₆		21 ₆	19 ₆
23									
24		11 ₆	19 ₂	13 ₆	11 ₆	*	*	13 ₆	*
25									
26									19 ₆
27		19 ₆			21 ₆	19 ₆		21 ₆	
28	25 ₁₀		13 ₁₀	25 ₁₀			13 ₁₀		
29		19 ₆			21 ₆	19 ₆		21 ₆	
30		19 ₆			21 ₆	19 ₆		21 ₆	19 ₆
31									
32		19 ₆			21 ₆	19 ₆		21 ₆	

B.

II. (concluded)

	A.									
	20	21	22	23	24	25	26	27	28	29
1	19 ₁	19 ₁	19 ₁	19 ₁	11 ₁	11 ₁	19 ₁	19 ₁	11 ₁	19 ₁
2	19 ₂	×	21 ₂		×	×			25 ₂	
3				19 ₂	25 ₂	24 ₂	21 ₂	19 ₂	25 ₂	21 ₂
4	19 ₂	19 ₁	19 ₁	19 ₂	11 ₁	11 ₁	19 ₁	19 ₂	11 ₁	19 ₁
5	19 ₂	19 ₁	19 ₁	19 ₆	11 ₆	3 ₁	21 ₆	19 ₆	3 ₁	21 ₆
6	19 ₆	×	21 ₆	19 ₁	3 ₁	11 ₆	19 ₁	19 ₂	3 ₁	19 ₁
7					25 ₂	25 ₂			*	
8	19 ₂	21 ₂	21 ₂		24 ₂	25 ₂			25 ₂	
9	19 ₂	21 ₂	21 ₂		11 ₆	3 ₁			3 ₁	
10	19 ₆	21 ₆	21 ₆		3 ₄	×			3 ₄	
11				19 ₂	25 ₂	24 ₂	21 ₂	19 ₂	25 ₂	21 ₂
12				19 ₆	25 ₁₀	3 ₄	21 ₆	19 ₆	3 ₄	21 ₆
13				19 ₂	3 ₁	11 ₆	21 ₂	19 ₂	3 ₁	21 ₂
14	*	19 ₁	19 ₁	19 ₆	11 ₆	3 ₁	21 ₆	19 ₆	3 ₁	21 ₆
15	19 ₆	21 ₆	21 ₆	19 ₂	3 ₁	11 ₆	19 ₁	*	3 ₁	19 ₁
B. 16	19 ₆	21 ₆	21 ₆	19 ₆	3 ₁	3 ₁	21 ₆	19 ₆	*	21 ₆
17					25 ₂	25 ₂			*	
18					25 ₁₀	3 ₄			3 ₄	
19					3 ₄	25 ₁₀			3 ₄	
20	19 ₂	21 ₂	21 ₂		11 ₆	3 ₁			3 ₁	
21	19 ₆	21 ₆	21 ₆		3 ₄	25 ₁₀			3 ₄	
22	19 ₆	21 ₆	21 ₆		3 ₄	3 ₄			*	
23				19 ₆	25 ₁₀	3 ₄	21 ₆	19 ₆	3 ₄	21 ₆
24				19 ₂	3 ₁	11 ₆	21 ₂	19 ₂	3 ₁	21 ₂
25		21 ₆	21 ₆	19 ₆	3 ₄	3 ₄	21 ₆	19 ₆	*	21 ₆
26	19 ₆	21 ₆	21 ₆	19 ₆	3 ₁	3 ₁	21 ₆	19 ₆	*	21 ₆
27					25 ₁₀	3 ₄			3 ₄	
28					3 ₄	25 ₁₀			3 ₄	
29					*	*			*	
30	19 ₆	21 ₆	21 ₆		3 ₄	3 ₄			*	
31				19 ₆	3 ₄	3 ₄	21 ₆	19 ₆	*	21 ₆
32					*	*			*	

For convenience the groups are divided into cases.

The double Table I gives all cases consistent with congruences (17), (21), (23) and (25). The results of the discussion are given in Table II. The cases in Table II left blank are inconsistent with congruences (22) and (24), and therefore have no groups corresponding to them.

Let $\kappa = \kappa_1 p^{k_2}$ where $dv[\kappa_1, p] = 1$ ($\kappa = a, \beta, c, g, \gamma, d, k, \alpha, \epsilon, e, j$).

In explanation of Table II the groups in cases marked $\boxed{r_s}$ are simply isomorphic with groups in $A_r B_s$.

The group G' is taken from the cases marked $\boxed{\times}$. The types are also selected from these cases.

The cases marked $\boxed{*}$ divide into two or more parts. Let

$$\begin{aligned}
a\epsilon - \alpha e + jk &= I_1, & a\epsilon - jk &= I_2, \\
a\delta(a - e) + 2I_1 &= I_3, & \alpha g - \beta j &= I_4, \\
\alpha\delta - \beta\epsilon &= I_5, & \epsilon g - \delta j &= I_6, \\
c\epsilon - e\gamma &= I_7, & \alpha e - jk &= I_8, \\
\delta e + \gamma j &= I_9, & \alpha\gamma + \delta k &= I_{10}.
\end{aligned}$$

The parts into which these groups divide, and the cases with which they are simply isomorphic, are given in Table III.

III.

$A_{1,2}B^*$	$dv[I_1, p] = p$	2_1	$dv[I_1, p] = 1$	2_4
A_3B^*	$dv[I_2, p] = p$	3_1	$dv[I_2, p] = 1$	3_4
A_4B^*	$dv[I_3, p] = p$	3_1	$dv[I_3, p] = 1$	3_4
$A_{12}B_{13}$	$dv[I_4, p] = p$	19_1	$dv[I_4, p] = 1$	19_2
$A_{14}B_{11}$	$dv[I_5, p] = p$	11_1	$dv[I_5, p] = 1$	24_2
$A_{15,18}B^*$	$dv[I_4, p] = p$	19_1	$dv[I_4, p] = 1$	21_2
$A_{16}B_{24}$	$dv[I_6, I_5, p] = p$	19_1	$dv[I_6, I_5, p] = 1$	19_2
$A_{20}B_{14}$	$dv[I_7, p] = p$	19_1	$dv[I_7, p] = 1$	19_2
$A_{24,25}B^*$	$dv[I_8, p] = p$	3_1	$dv[I_8, p] = 1$	3_4
$A_{27}B_{15}$	$dv[I_6, p] = p$	19_1	$dv[I_6, p] = 1$	19_2
$A_{29}B_{7,17}$	$dv[I_{10}, p] = p$	24_2	$dv[I_{10}, p] = 1$	25_2
$A_{29}B_{16,26}$	$dv[I_9, p] = p$	11_6	$dv[I_9, p] = 1$	3_1
$A_{29}B_{22,25,30,31}$	$dv[I_9, p] = p$	25_{10}	$dv[I_9, p] = 1$	3_4
$A_{29}B_{29,32}$	$dv[I_8, I_9, p] = p$	11_6	$[I_8, p] = p, [I_9, p] = 1$	3_1
$A_{29}B_{29,32}$	$[I_8, p] = 1, [I_9, p] = p$	25_{10}	$[I_8, p] = 1, [I_9, p] = 1$	3_4

8. *Types.* The types for this class are given by equations (30) where the constants have the values given in Table IV.

IV.

	a	β	c	g	γ	δ	k	α	ϵ	e	j
1_1	0	0	0	0	0	0	0	0	0	0	0
2_1	1	0	0	0	0	0	0	0	0	0	0
3_1	κ	1	0	0	0	0	0	0	0	0	0
11_1	0	1	0	0	0	0	0	0	0	0	0
* 13_1	0	1	0	0	1	0	0	0	0	0	0
19_1	0	0	1	0	0	0	0	0	0	0	0
1_2	0	0	0	0	0	0	1	0	0	0	0
* 13_2	0	1	0	0	1	0	1	0	0	0	0
19_2	0	0	1	0	0	0	1	0	0	0	0
* 21_2	0	0	1	0	0	1	1	0	0	0	0
24_2	0	0	0	0	1	0	1	0	0	0	0
25_2	0	0	0	0	0	1	1	0	0	0	0
2_4	1	0	0	0	0	0	0	0	1	0	0
3_4	κ	1	0	0	0	0	0	0	1	0	0
11_6	0	1	0	0	0	0	0	0	0	0	1
* 13_6	0	1	0	0	1	0	0	0	0	0	1

$\kappa = 1$, and a non-residue (mod p).

*For $p = 3$ these groups are isomorphic in Class II.

A detailed analysis of congruences (32) for several cases is given below as a general illustration of the methods used.

A_3B_1 .

The special forms of the congruences for this case are

$$\beta'xz' \equiv 0 \pmod{p}, \quad \text{(II)}$$

$$a'(yz' - y'z) \equiv kx \pmod{p}, \quad \text{(III)}$$

$$\beta v' \equiv 0, \quad \beta z' \equiv 0, \quad \beta y' \equiv \beta'xz'' \pmod{p}, \quad \text{(IV),(V),(VI)}$$

$$a'(yz'' - y''z) + a'\beta'x\binom{z''}{2} \equiv \alpha x + \beta x' + a'\beta y'z \pmod{p}, \quad \text{(VII)}$$

$$a'(y'z'' - y''z') \equiv ax \pmod{p}, \quad \text{(X)}$$

$$\gamma v'' + \delta v' \equiv 0 \pmod{p}, \quad \text{(XI)}$$

$$\gamma z'' + \delta z' \equiv 0 \pmod{p}, \quad \text{(XII)}$$

$$\gamma y'' + \delta y' \equiv \beta'xz'' \pmod{p}, \quad \text{(XIII)}$$

$$a'(yz''' - y'''z) + a'\beta'x\binom{z'''}{2} \equiv \epsilon x + \gamma x'' + \delta x + a'\delta y'z + a'\gamma y''z + a'\binom{\gamma}{2}y''z'' \pmod{p}, \quad \text{(XIV)}$$

$$cv'' \equiv 0, \quad cz'' \equiv 0, \quad cy'' \equiv 0 \pmod{p}, \quad \text{(XV),(XVI),(XVII)}$$

$$a'(y'z''' - y'''z') \equiv ex \pmod{p}, \quad \text{(XVIII)}$$

$$\begin{aligned}
gv' \equiv 0, \quad gz' \equiv 0, \quad gy' \equiv 0 \pmod{p}, & \quad (\text{XIX}), (\text{XX}), (\text{XXI}) \\
a'(y''z''' - y'''z'') \equiv jx \pmod{p}, & \quad (\text{XXII})
\end{aligned}$$

From (II) $z' \equiv 0 \pmod{p}$.

The conditions of isomorphism give

$$\Delta \equiv \begin{vmatrix} v' & v'' & v''' \\ y' & y'' & y''' \\ z' & z'' & z''' \end{vmatrix} \not\equiv 0 \pmod{p}.$$

Multiply (IV), (V), (VI) by γ and reduce by (XII), $\beta\gamma v' \equiv 0$, $\beta\gamma z' \equiv 0$, $\beta\gamma y' \equiv 0 \pmod{p}$. Since $\Delta \not\equiv 0 \pmod{p}$, one at least of the quantities, v' , z' or y' is $\not\equiv 0 \pmod{p}$ and $\beta\gamma \equiv 0 \pmod{p}$.

From (XV), (XVI) and (XVII) $c \equiv 0 \pmod{p}$, and from (XIX), (XX) and (XXI) $g \equiv 0 \pmod{p}$.

From (IV), (V), (VI) and (X) if $a \equiv 0$, then $\beta \equiv 0$ and if $a \not\equiv 0$, then $\beta \not\equiv 0 \pmod{p}$.

At least one of the three quantities β , γ or δ is $\not\equiv 0 \pmod{p}$ and one, at least, of a , e or j is $\not\equiv 0 \pmod{p}$.

A_3 : Since $z''' \equiv 0 \pmod{p}$, (XVIII) gives $e \equiv 0$. Elimination between (III), (X), (XIV) and (XXII) gives $a\epsilon - kj \equiv 0 \pmod{p}$. Elimination between (VI) and (X) gives $a'\beta'z''^2 \equiv a\beta \pmod{p}$ and $a\beta$ is a quadratic residue or non-residue according as $a'\beta'$ is or is not, and there are two types for this case.

A_4 : Since y' and z'' are $\not\equiv 0 \pmod{p}$, $e \not\equiv 0 \pmod{p}$. Elimination between (VI), (X), (XIII) and (XVIII) gives $a\delta - \beta e \equiv 0 \pmod{p}$.

This is a special form of (24).

Elimination between (III), (VII), (X), (XIII), (XIV), (XVIII) and (XXII) gives

$$2jk + a\delta(a - e) + 2(a\epsilon - \alpha e) \equiv 0 \pmod{p}.$$

A_{24} : Since from (XI), (XII) and (XIII) y'' and $z''' \not\equiv 0 \pmod{p}$, and $z'' \equiv v'' \equiv 0 \pmod{p}$, (xxii) gives $j \not\equiv 0 \pmod{p}$.

Elimination between (III), (X), (XVIII) and (XXII) gives

$$\alpha e - jk \equiv 0 \pmod{p}.$$

A_{25} : (XI), (XII) and (XIII) give $v' \equiv z' \equiv 0$ and $y', z''' \not\equiv 0 \pmod{p}$ and this with (XVIII) gives $e \not\equiv 0$.

Elimination between (III), (VII), (XVIII) and (XXII) gives

$$\alpha e - jk \equiv 0 \pmod{p}.$$

A_{28} : Since $a \equiv 0$ then e or $j \not\equiv 0 \pmod{p}$.

Elimination between (III), (VII), (XVIII) and (XXII) gives

$$\alpha e - jk \equiv 0 \pmod{p}.$$

Multiply (XIII) by $a'z'''$ and reduce

$$\delta e + \gamma j \equiv a' \beta' z'''^2 \not\equiv 0 \pmod{p}.$$

$$A_{11}B_1.$$

The special forms of the congruences for this case are

$$\beta'xz' \equiv 0 \pmod{p}, \quad (\text{II})$$

$$kx \equiv 0 \pmod{p}, \quad (\text{III})$$

$$\beta v' \equiv \beta z' \equiv 0, \quad \beta y' \equiv \beta'xz'', \quad (\text{IV}),(\text{V}),(\text{VI})$$

$$\alpha x + \beta x' \equiv 0 \pmod{p}, \quad (\text{VII})$$

$$ax \equiv 0 \pmod{p}, \quad (\text{X})$$

$$\gamma v'' + \delta v \equiv 0 \pmod{p}, \quad (\text{XI})$$

$$\gamma z'' + \delta z \equiv 0 \pmod{p}, \quad (\text{XII})$$

$$\gamma y'' + \delta y \equiv \beta'xz'''' \pmod{p}, \quad (\text{XIII})$$

$$\epsilon x + \gamma x'' + \delta x' \equiv 0 \pmod{p}, \quad (\text{XIV})$$

$$cv'' \equiv cz'' \equiv cy'' \equiv 0 \pmod{p}, \quad (\text{XV}),(\text{XVI}),(\text{XVII})$$

$$ex \equiv 0 \pmod{p}, \quad (\text{XVIII})$$

$$gv' \equiv gz' \equiv gy' \equiv 0 \pmod{p}, \quad (\text{XIX}),(\text{XX}),(\text{XXI})$$

$$jx \equiv 0 \pmod{p}, \quad (\text{XXII})$$

(II) gives $z' = 0$, (III) gives $k \equiv 0$, (X) gives $a \equiv 0$, (XV), (XVI), (XVII) give $c \equiv 0$ ($\Delta \not\equiv 0$), (XVIII) gives $e \equiv 0$, (XIX), (XX), (XXI) give $g \equiv 0$, (XXII) gives $j \equiv 0$. One of the two quantities z'' or $z'''' \not\equiv 0 \pmod{p}$, and by (VI) and (XIII) one of the three quantities β , γ or δ is $\not\equiv 0$.

A_{11} : (XIV) gives $\epsilon \equiv 0 \pmod{p}$. Multiplying (IV), (V), (VI) by γ gives, by (XII), $\beta\gamma v' \equiv \beta\gamma z' \equiv \beta\gamma y' \equiv 0 \pmod{p}$, and $\beta\gamma \equiv 0 \pmod{p}$.

A_{14} : Elimination between (VII) and (XIV) gives $\alpha\delta - \beta\epsilon \equiv 0 \pmod{p}$.

A_{24} : (VII) gives $\alpha \equiv 0 \pmod{p}$, (XIV) $\epsilon \equiv 0$ or $\not\equiv 0 \pmod{p}$.

A_{25} : (VII) gives $\alpha \equiv 0 \pmod{p}$, (XIV) $\epsilon \equiv$ or $\not\equiv 0 \pmod{p}$.

A_{28} : (VII) gives $\alpha \equiv 0 \pmod{p}$, (XIV) $\epsilon \equiv$ or $\not\equiv 0 \pmod{p}$.

$A_{19}B_1$.

The special forms of the congruences for this case are

$$\begin{aligned}
c'(yv' - y'v) &\equiv 0 \pmod{p}, & (I) \\
kx &\equiv 0 \pmod{p}, & (III) \\
\beta v \equiv 0, \quad \beta z &\equiv c'(yv'' - y''v), \quad \beta y' \equiv 0 \pmod{p}, & (IV),(V),(VI) \\
\alpha x + \beta x' &\equiv 0 \pmod{p}, & (VII) \\
c'(y'v'' - y''v') &\equiv 0 \pmod{p}, & (VIII) \\
ax &\equiv 0 \pmod{p}, & (X) \\
\gamma v'' + \delta v' &\equiv 0 \pmod{p}, & (XI) \\
\gamma z'' + \delta z' + c'\gamma\delta y''v + c' \binom{\delta}{2} v'y' + c' \binom{\gamma}{2} v''y'' &\equiv c'(yv''' - y'''v) \pmod{p}, & (XII) \\
\gamma y'' + \delta y' &\equiv 0 \pmod{p}, & (XIII) \\
\epsilon x + \gamma x'' + \delta x' &\equiv 0 \pmod{p}, & (XIV) \\
cv'' \equiv 0, \quad cz'' &\equiv c'(y'v''' - y'''v'), \quad cy'' \equiv 0 \pmod{p}, & (XV),(XVI),(XVII) \\
ex + cx'' &\equiv 0 \pmod{p}, & (XVIII) \\
gv' \equiv 0, \quad gz' &\equiv c'(y''v''' - y'''v''), \quad gy' \equiv 0 \pmod{p}, & (XIX),(XX),(XXI) \\
jx + gx' &\equiv 0 \pmod{p}. & (XXII)
\end{aligned}$$

(III) gives $k \equiv 0$, (X) gives $a \equiv 0$.

Since $dv[(y'v''' - y'''v'), (y''v''' - y'''v''), p] = 1$ then $dv[c, g, p] = 1$.

If $c \not\equiv 0$, $v'' \equiv y'' \equiv 0 \pmod{p}$ and therefore $g \equiv 0 \pmod{p}$ and if $g \not\equiv 0$, then $c \equiv 0 \pmod{p}$.

A_{12} : (XVIII) gives $e \equiv 0 \pmod{p}$. Elimination between (VII) and (XXII) gives $\alpha g - \beta j \equiv 0 \pmod{p}$, (XIV) gives $\epsilon \equiv 0 \pmod{p}$.

A_{15} : (XVIII) gives $e \equiv 0 \pmod{p}$. Elimination between (VII) and (XXII) gives $\alpha g - \beta j \equiv 0 \pmod{p}$, (XIV) gives $\epsilon \equiv 0$ or $\not\equiv 0 \pmod{p}$.

A_{16} : (XVIII) gives $e \equiv 0$. Elimination between (XIV) and (XXII) gives $\epsilon g - \delta j \equiv 0 \pmod{p}$, between (VII) and (XIV) gives $\alpha\delta - \beta\epsilon \equiv 0$.

A_{18} : (XVIII) gives $e \equiv 0 \pmod{p}$. Elimination between (VII) and (XXII) gives $\alpha g - \beta j \equiv 0 \pmod{p}$, (XIV) gives $\epsilon \equiv 0$ or $\not\equiv 0 \pmod{p}$.

A_{19} : (VII) gives $\alpha \equiv 0 \pmod{p}$, (XIV) gives $\epsilon \equiv 0$, (XXII) gives $j \equiv 0 \pmod{p}$, (XVIII) gives $e \equiv 0$ or $\not\equiv 0 \pmod{p}$.

A_{20} : (VII) gives $\alpha \equiv 0$, (XXII) gives $j \equiv 0$. Elimination between (XIV) and (XVIII) gives $\epsilon c - e\gamma \equiv 0 \pmod{p}$.

A_{21} : (VII) gives $\alpha \equiv 0$, (XIV) gives $\epsilon \equiv 0$ or $\not\equiv 0 \pmod{p}$, (XVIII) gives $e \equiv 0$, or $\not\equiv 0$, and (XXII) gives $j \equiv 0 \pmod{p}$.

A_{22} : (VII) gives $\alpha \equiv 0$, (XIV) gives $\epsilon \equiv 0$ or $\not\equiv 0$, (XVIII) gives $\epsilon \equiv 0$ or $\not\equiv 0$, (XXII) gives $j \equiv 0 \pmod{p}$.

A_{23} : (VII) gives $\alpha \equiv 0$, (XIV) gives $\epsilon \equiv 0$, (XVIII) gives $\epsilon \equiv 0$, (XXII) gives $j \equiv 0$ or $\not\equiv 0 \pmod{p}$.

A_{26} : (VII) $\alpha \equiv 0$, (XIV) $\epsilon \equiv 0$ or $\not\equiv 0$, (XVIII) $\epsilon \equiv 0$, (XXII) $j \equiv 0$ or $\not\equiv 0 \pmod{p}$.

A_{27} : (VII) $\alpha \equiv 0$, (XIV) $\epsilon \equiv 0$ or $\not\equiv 0$, (XVIII) $\epsilon \equiv 0$, (XXII) $j \equiv 0$ or $\not\equiv 0 \pmod{p}$. Elimination between (XIV) and (XXII) gives $\epsilon g - \delta j \equiv 0 \pmod{p}$.

A_{29} : (VII) $\alpha \equiv 0$, (XIV) $\epsilon \equiv 0$ or $\not\equiv 0$, (XVIII) $\epsilon \equiv 0$, (XXII) $j \equiv 0$ or $\not\equiv 0 \pmod{p}$.

End of the Project Gutenberg EBook of Groups of Order p^m Which Contain
Cyclic Subgroups of Order $p^{(m-3)}$, by Lewis Irving Neikirk

*** END OF THIS PROJECT GUTENBERG EBOOK GROUPS OF ORDER P^M WHICH ***

***** This file should be named 9930-pdf.pdf or 9930-pdf.zip *****
This and all associated files of various formats will be found in:
<http://www.gutenberg.org/9/9/3/9930/>

Produced by Cornell University, Joshua Hutchinson, Lee
Chew-Hung, John Hagerson, and the Online Distributed
Proofreading Team

Updated editions will replace the previous one--the old editions
will be renamed.

Creating the works from public domain print editions means that no
one owns a United States copyright in these works, so the Foundation
(and you!) can copy and distribute it in the United States without
permission and without paying copyright royalties. Special rules,
set forth in the General Terms of Use part of this license, apply to
copying and distributing Project Gutenberg-tm electronic works to
protect the PROJECT GUTENBERG-tm concept and trademark. Project
Gutenberg is a registered trademark, and may not be used if you
charge for the eBooks, unless you receive specific permission. If you
do not charge anything for copies of this eBook, complying with the
rules is very easy. You may use this eBook for nearly any purpose
such as creation of derivative works, reports, performances and
research. They may be modified and printed and given away--you may do
practically ANYTHING with public domain eBooks. Redistribution is
subject to the trademark license, especially commercial
redistribution.

*** START: FULL LICENSE ***

THE FULL PROJECT GUTENBERG LICENSE
PLEASE READ THIS BEFORE YOU DISTRIBUTE OR USE THIS WORK

To protect the Project Gutenberg-tm mission of promoting the free
distribution of electronic works, by using or distributing this work
(or any other work associated in any way with the phrase "Project
Gutenberg"), you agree to comply with all the terms of the Full Project
Gutenberg-tm License available with this file or online at
www.gutenberg.org/license.

Section 1. General Terms of Use and Redistributing Project Gutenberg-tm
electronic works

1.A. By reading or using any part of this Project Gutenberg-tm electronic work, you indicate that you have read, understand, agree to and accept all the terms of this license and intellectual property (trademark/copyright) agreement. If you do not agree to abide by all the terms of this agreement, you must cease using and return or destroy all copies of Project Gutenberg-tm electronic works in your possession. If you paid a fee for obtaining a copy of or access to a Project Gutenberg-tm electronic work and you do not agree to be bound by the terms of this agreement, you may obtain a refund from the person or entity to whom you paid the fee as set forth in paragraph 1.E.8.

1.B. "Project Gutenberg" is a registered trademark. It may only be used on or associated in any way with an electronic work by people who agree to be bound by the terms of this agreement. There are a few things that you can do with most Project Gutenberg-tm electronic works even without complying with the full terms of this agreement. See paragraph 1.C below. There are a lot of things you can do with Project Gutenberg-tm electronic works if you follow the terms of this agreement and help preserve free future access to Project Gutenberg-tm electronic works. See paragraph 1.E below.

1.C. The Project Gutenberg Literary Archive Foundation ("the Foundation" or PGLAF), owns a compilation copyright in the collection of Project Gutenberg-tm electronic works. Nearly all the individual works in the collection are in the public domain in the United States. If an individual work is in the public domain in the United States and you are located in the United States, we do not claim a right to prevent you from copying, distributing, performing, displaying or creating derivative works based on the work as long as all references to Project Gutenberg are removed. Of course, we hope that you will support the Project Gutenberg-tm mission of promoting free access to electronic works by freely sharing Project Gutenberg-tm works in compliance with the terms of this agreement for keeping the Project Gutenberg-tm name associated with the work. You can easily comply with the terms of this agreement by keeping this work in the same format with its attached full Project Gutenberg-tm License when you share it without charge with others.

1.D. The copyright laws of the place where you are located also govern what you can do with this work. Copyright laws in most countries are in a constant state of change. If you are outside the United States, check the laws of your country in addition to the terms of this agreement before downloading, copying, displaying, performing, distributing or creating derivative works based on this work or any other Project Gutenberg-tm work. The Foundation makes no representations concerning the copyright status of any work in any country outside the United States.

1.E. Unless you have removed all references to Project Gutenberg:

1.E.1. The following sentence, with active links to, or other immediate access to, the full Project Gutenberg-tm License must appear prominently whenever any copy of a Project Gutenberg-tm work (any work on which the

phrase "Project Gutenberg" appears, or with which the phrase "Project Gutenberg" is associated) is accessed, displayed, performed, viewed, copied or distributed:

This eBook is for the use of anyone anywhere at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at www.gutenberg.org

1.E.2. If an individual Project Gutenberg-tm electronic work is derived from the public domain (does not contain a notice indicating that it is posted with permission of the copyright holder), the work can be copied and distributed to anyone in the United States without paying any fees or charges. If you are redistributing or providing access to a work with the phrase "Project Gutenberg" associated with or appearing on the work, you must comply either with the requirements of paragraphs 1.E.1 through 1.E.7 or obtain permission for the use of the work and the Project Gutenberg-tm trademark as set forth in paragraphs 1.E.8 or 1.E.9.

1.E.3. If an individual Project Gutenberg-tm electronic work is posted with the permission of the copyright holder, your use and distribution must comply with both paragraphs 1.E.1 through 1.E.7 and any additional terms imposed by the copyright holder. Additional terms will be linked to the Project Gutenberg-tm License for all works posted with the permission of the copyright holder found at the beginning of this work.

1.E.4. Do not unlink or detach or remove the full Project Gutenberg-tm License terms from this work, or any files containing a part of this work or any other work associated with Project Gutenberg-tm.

1.E.5. Do not copy, display, perform, distribute or redistribute this electronic work, or any part of this electronic work, without prominently displaying the sentence set forth in paragraph 1.E.1 with active links or immediate access to the full terms of the Project Gutenberg-tm License.

1.E.6. You may convert to and distribute this work in any binary, compressed, marked up, nonproprietary or proprietary form, including any word processing or hypertext form. However, if you provide access to or distribute copies of a Project Gutenberg-tm work in a format other than "Plain Vanilla ASCII" or other format used in the official version posted on the official Project Gutenberg-tm web site (www.gutenberg.org), you must, at no additional cost, fee or expense to the user, provide a copy, a means of exporting a copy, or a means of obtaining a copy upon request, of the work in its original "Plain Vanilla ASCII" or other form. Any alternate format must include the full Project Gutenberg-tm License as specified in paragraph 1.E.1.

1.E.7. Do not charge a fee for access to, viewing, displaying, performing, copying or distributing any Project Gutenberg-tm works unless you comply with paragraph 1.E.8 or 1.E.9.

1.E.8. You may charge a reasonable fee for copies of or providing access to or distributing Project Gutenberg-tm electronic works provided that

- You pay a royalty fee of 20% of the gross profits you derive from the use of Project Gutenberg-tm works calculated using the method you already use to calculate your applicable taxes. The fee is owed to the owner of the Project Gutenberg-tm trademark, but he has agreed to donate royalties under this paragraph to the Project Gutenberg Literary Archive Foundation. Royalty payments must be paid within 60 days following each date on which you prepare (or are legally required to prepare) your periodic tax returns. Royalty payments should be clearly marked as such and sent to the Project Gutenberg Literary Archive Foundation at the address specified in Section 4, "Information about donations to the Project Gutenberg Literary Archive Foundation."
- You provide a full refund of any money paid by a user who notifies you in writing (or by e-mail) within 30 days of receipt that s/he does not agree to the terms of the full Project Gutenberg-tm License. You must require such a user to return or destroy all copies of the works possessed in a physical medium and discontinue all use of and all access to other copies of Project Gutenberg-tm works.
- You provide, in accordance with paragraph 1.F.3, a full refund of any money paid for a work or a replacement copy, if a defect in the electronic work is discovered and reported to you within 90 days of receipt of the work.
- You comply with all other terms of this agreement for free distribution of Project Gutenberg-tm works.

1.E.9. If you wish to charge a fee or distribute a Project Gutenberg-tm electronic work or group of works on different terms than are set forth in this agreement, you must obtain permission in writing from both the Project Gutenberg Literary Archive Foundation and Michael Hart, the owner of the Project Gutenberg-tm trademark. Contact the Foundation as set forth in Section 3 below.

1.F.

1.F.1. Project Gutenberg volunteers and employees expend considerable effort to identify, do copyright research on, transcribe and proofread public domain works in creating the Project Gutenberg-tm collection. Despite these efforts, Project Gutenberg-tm electronic works, and the medium on which they may be stored, may contain "Defects," such as, but not limited to, incomplete, inaccurate or corrupt data, transcription errors, a copyright or other intellectual property infringement, a defective or damaged disk or other medium, a computer virus, or computer codes that damage or cannot be read by

your equipment.

1.F.2. LIMITED WARRANTY, DISCLAIMER OF DAMAGES - Except for the "Right of Replacement or Refund" described in paragraph 1.F.3, the Project Gutenberg Literary Archive Foundation, the owner of the Project Gutenberg-tm trademark, and any other party distributing a Project Gutenberg-tm electronic work under this agreement, disclaim all liability to you for damages, costs and expenses, including legal fees. YOU AGREE THAT YOU HAVE NO REMEDIES FOR NEGLIGENCE, STRICT LIABILITY, BREACH OF WARRANTY OR BREACH OF CONTRACT EXCEPT THOSE PROVIDED IN PARAGRAPH 1.F.3. YOU AGREE THAT THE FOUNDATION, THE TRADEMARK OWNER, AND ANY DISTRIBUTOR UNDER THIS AGREEMENT WILL NOT BE LIABLE TO YOU FOR ACTUAL, DIRECT, INDIRECT, CONSEQUENTIAL, PUNITIVE OR INCIDENTAL DAMAGES EVEN IF YOU GIVE NOTICE OF THE POSSIBILITY OF SUCH DAMAGE.

1.F.3. LIMITED RIGHT OF REPLACEMENT OR REFUND - If you discover a defect in this electronic work within 90 days of receiving it, you can receive a refund of the money (if any) you paid for it by sending a written explanation to the person you received the work from. If you received the work on a physical medium, you must return the medium with your written explanation. The person or entity that provided you with the defective work may elect to provide a replacement copy in lieu of a refund. If you received the work electronically, the person or entity providing it to you may choose to give you a second opportunity to receive the work electronically in lieu of a refund. If the second copy is also defective, you may demand a refund in writing without further opportunities to fix the problem.

1.F.4. Except for the limited right of replacement or refund set forth in paragraph 1.F.3, this work is provided to you 'AS-IS', WITH NO OTHER WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR ANY PURPOSE.

1.F.5. Some states do not allow disclaimers of certain implied warranties or the exclusion or limitation of certain types of damages. If any disclaimer or limitation set forth in this agreement violates the law of the state applicable to this agreement, the agreement shall be interpreted to make the maximum disclaimer or limitation permitted by the applicable state law. The invalidity or unenforceability of any provision of this agreement shall not void the remaining provisions.

1.F.6. INDEMNITY - You agree to indemnify and hold the Foundation, the trademark owner, any agent or employee of the Foundation, anyone providing copies of Project Gutenberg-tm electronic works in accordance with this agreement, and any volunteers associated with the production, promotion and distribution of Project Gutenberg-tm electronic works, harmless from all liability, costs and expenses, including legal fees, that arise directly or indirectly from any of the following which you do or cause to occur: (a) distribution of this or any Project Gutenberg-tm work, (b) alteration, modification, or additions or deletions to any Project Gutenberg-tm work, and (c) any Defect you cause.

Section 2. Information about the Mission of Project Gutenberg-tm

Project Gutenberg-tm is synonymous with the free distribution of electronic works in formats readable by the widest variety of computers including obsolete, old, middle-aged and new computers. It exists because of the efforts of hundreds of volunteers and donations from people in all walks of life.

Volunteers and financial support to provide volunteers with the assistance they need are critical to reaching Project Gutenberg-tm's goals and ensuring that the Project Gutenberg-tm collection will remain freely available for generations to come. In 2001, the Project Gutenberg Literary Archive Foundation was created to provide a secure and permanent future for Project Gutenberg-tm and future generations. To learn more about the Project Gutenberg Literary Archive Foundation and how your efforts and donations can help, see Sections 3 and 4 and the Foundation information page at www.gutenberg.org

Section 3. Information about the Project Gutenberg Literary Archive Foundation

The Project Gutenberg Literary Archive Foundation is a non profit 501(c)(3) educational corporation organized under the laws of the state of Mississippi and granted tax exempt status by the Internal Revenue Service. The Foundation's EIN or federal tax identification number is 64-6221541. Contributions to the Project Gutenberg Literary Archive Foundation are tax deductible to the full extent permitted by U.S. federal laws and your state's laws.

The Foundation's principal office is located at 4557 Melan Dr. S. Fairbanks, AK, 99712., but its volunteers and employees are scattered throughout numerous locations. Its business office is located at 809 North 1500 West, Salt Lake City, UT 84116, (801) 596-1887. Email contact links and up to date contact information can be found at the Foundation's web site and official page at www.gutenberg.org/contact

For additional contact information:

Dr. Gregory B. Newby
Chief Executive and Director
gnewby@pglaf.org

Section 4. Information about Donations to the Project Gutenberg Literary Archive Foundation

Project Gutenberg-tm depends upon and cannot survive without wide spread public support and donations to carry out its mission of increasing the number of public domain and licensed works that can be freely distributed in machine readable form accessible by the widest array of equipment including outdated equipment. Many small donations

(\$1 to \$5,000) are particularly important to maintaining tax exempt status with the IRS.

The Foundation is committed to complying with the laws regulating charities and charitable donations in all 50 states of the United States. Compliance requirements are not uniform and it takes a considerable effort, much paperwork and many fees to meet and keep up with these requirements. We do not solicit donations in locations where we have not received written confirmation of compliance. To SEND DONATIONS or determine the status of compliance for any particular state visit www.gutenberg.org/donate

While we cannot and do not solicit contributions from states where we have not met the solicitation requirements, we know of no prohibition against accepting unsolicited donations from donors in such states who approach us with offers to donate.

International donations are gratefully accepted, but we cannot make any statements concerning tax treatment of donations received from outside the United States. U.S. laws alone swamp our small staff.

Please check the Project Gutenberg Web pages for current donation methods and addresses. Donations are accepted in a number of other ways including checks, online payments and credit card donations. To donate, please visit: www.gutenberg.org/donate

Section 5. General Information About Project Gutenberg-tm electronic works.

Professor Michael S. Hart was the originator of the Project Gutenberg-tm concept of a library of electronic works that could be freely shared with anyone. For forty years, he produced and distributed Project Gutenberg-tm eBooks with only a loose network of volunteer support.

Project Gutenberg-tm eBooks are often created from several printed editions, all of which are confirmed as Public Domain in the U.S. unless a copyright notice is included. Thus, we do not necessarily keep eBooks in compliance with any particular paper edition.

Most people start at our Web site which has the main PG search facility:

www.gutenberg.org

This Web site includes information about Project Gutenberg-tm, including how to make donations to the Project Gutenberg Literary Archive Foundation, how to help produce our new eBooks, and how to subscribe to our email newsletter to hear about new eBooks.